



# Ameliorative Effect of Rosemary Extract (*Rosmarinus Officinalis* L.) Against Ethion Bound Residues Induced Alterations in Experimental Animals

Abdel-Gawad H<sup>1\*</sup>, Taha H<sup>1</sup> and Sayed Aly MA<sup>2</sup>

<sup>1</sup>Applied Organic Chemistry Department, Chemical Industries Research Division, National Research Centre, Egypt

<sup>2</sup>Department of Animal Reproduction and Artificial Insemination, Veterinary Research Division, National Research Centre, National Research Centre, Egypt

**\*Corresponding author:** Hassan Abdel-Gawad, Applied Organic Chemistry Department, Chemical Industries Research Division, National Research Centre, 33 EL Buhouth St, Dokki, Giza 12622, Egypt, Email: abdelgawadhassan@hotmail.com

**Received Date:** February 05, 2020; **Published Date:** February 27, 2020

## Abstract

Fatal poisoning is widespread upon human exposure to toxic agents such as pesticides, heavy metals and environmental pollutants. In this regards, the application of the natural dietary agents as antidotes has engrossed a substantial attention. One of the ancient known traditional medicines and spices with an arsenal of metabolites of several reported health benefits is Rosemary (*Rosmarinus officinalis* L.). The current study was implemented to demonstrate the protective and nutritional impact of rosemary leaves extract (1% w/w) in experimental rats fed on maize cake containing ethion bound residues (1.6 ppm). Feeding animals with treated cake containing ethion bound residues caused elevation in serum alanine aminotransferase, aspartate aminotransferase, and alkaline phosphatase activities ( $P < 0.01$ ,  $P < 0.001$  and  $P < 0.01$ ), respectively. Whereas total protein and albumin were significantly decreased ( $P < 0.01$  and  $P < 0.05$ ), respectively.

As well as, concomitant elevation in serum creatinine and urea levels ( $P < 0.01$ ). Also, a significant ( $P < 0.01$ ) increase in cholesterol, triglyceride and low density lipoprotein and a significant ( $P < 0.001$ ) decrease in high density lipoprotein content as compared to the control were obtained. In addition a decrease in acetyl cholinesterase, catalase, and glutathione-S-transferase activity was observed as compared with controls. Furthermore, acute ethion bound residues intoxication caused histopathological changes in liver, kidney and brain. Administration of rosemary shows better effect in the activity of liver enzymes, kidney function markers, lipids contents as well as, protein profiles and antioxidant status. Rosemary also exhibited some improvement in the histological architecture of liver, kidney and brain.

**Keywords:** Ethion bound residues; Rosemary; Hepatotoxicity; Nephrotoxicity; Lipid profile; Rat

**Abbreviations:** Ops: Organophosphorus; CBs: TC: Total Cholesterol; TG: Triglycerides; HDL: High Density Carbamates; PYs: Pyrethroids; OPIs: Organophosphorus insecticides; ALT: Alanine Aminotransferase; AST: Aspartate Aminotransferase; ALP: Alkaline Phosphatase; Lipoprotein; LDL: Low Density Lipoprotein; GST: Glutathione-S-Transferase.

## Introduction

Pesticides and agrochemicals, in general, became an important component of worldwide agriculture systems during the last century, allowing for a noticeable increase in crop yields and food production. Exposure to either natural or synthetic chemicals represents a worldwide public health problem. Humans or animals could be exposed to pesticides in air, soil, food, fruits, vegetables, or even pharmaceutical and cosmetic products via inhalation, ingestion, or direct contact. Among the many classes of pesticides to which humans are exposed to via the diet are the organophosphorus (OPs), carbamates (CBs) and pyrethroids (PYs), which mechanisms of actions to their target organism also occur in human and other mammals. At the same time, during the last decades we realized that agrochemical residues did spread in the environment, causing significant contamination of terrestrial ecosystems and poisoning human foods [1-5].

Maize crop is one of the food crops that have several uses, whether as a food for man or as animal feed due to its high nutrition value. The use of pesticides can result in the accumulation of their residues in seeds, oil and cake. The residues in cake reach the milk through lactating animals and eventually into human body. Since consumers could be at risk from the pesticide residues which may be present in cake, therefore it is important to develop a procedure for the removal of pesticide residue from cakes. Ethion (O,O,O',O'-tetraethyl-S,S'-methylene-bis phosphorodithioate) is one of the widely used organophosphorus insecticides (OPs), which have been identified as contaminants in many component of the global ecosystem. It is used to kill aphids, mites, thrips, leafhoppers and maggots on crops, ornamentals and animals. Ethion is converted in the liver to its active oxygenated analog, ethion mono-oxon, via desulfuration by cytochrome P-450 enzymes [6-10].


Natural herbs are widely consumed by humans on a daily basis; these natural products have many biologic and pharmacologic properties. Rosemary (*Rosmarinus officinalis* L.) (RM) is one of the most widely used and well- documented medicinal plants in the world, it is used as flavouring agent for food, beverages, and cosmetics preparations. The phytochemical screening of this plant has revealed its richness in beneficial active molecules such as polyphenolic compounds. For this reason, rosemary extracts exhibit many beneficial health effects as antiviral, radioprotective, anti-microbial, neuroprotective, anticancer, anti-diarrheal and anti-inflammatory properties, hypolipidaemic agent, hepatoprotective (and nephroprotective [11-23]. Rosemary (*Rosmarinus officinalis* L.), contains antioxidants such as rosmarinic acid, diterpenoids like carnosic acid, carnosol, rosmanol, and epirosmanol as well as tocopherols and carotenoids. The antioxidant activities of Rosemary (*Rosmarinus officinalis*

L.) extract have been associated with phenolic diterpenes that scavenge singlet oxygen, hydroxyl radicals and lipid peroxy radicals [24-27]. The protective role of rosemary extract against ethion bound residue - induced hepatic, nephro, and neurotoxicity is not fully explored. Therefore, the present study was planned to evaluate protective effects of rosemary aqueous extract against ethion bound residue - induced toxicity by examining hematological alterations, histopathological changes and oxidative stress.

## Materials and Methods

### <sup>14</sup>C-Labeled ethion

<sup>14</sup>C-ethion labeled in the ethoxy groups (Figure 1) was prepared in a single vessel reaction using <sup>14</sup>C-ethyl alcohol (Sp. Act. 37 MBq, Amersham, England) according to Abdel-Gawad et al [28]. <sup>14</sup>C-Ethion had a specific activity of 7.4 MBq/g and the radiometric purity was 98 %. The R<sub>f</sub> values of ethion in different solvent systems (Toluene: Xylene 20: 20; Dioxane: Xylene: Petroleum ether 10: 20: 20 and n-Hexane: Ethyl acetate 98: 2) are 0.75, 0.8 and 0.76, respectively. The absorbance of ethion was measured by UV spectrophotometer (JASCO) at λ=290 nm.



### Plant material and experimental design

Seeds of maize were cultivated under normal field conditions in a controlled, isolated field area as practiced in the field. Shortly at blooming stage, leaves of plants were treated twice, 15 days apart, with <sup>14</sup>C-ethion at the dose of 4 mg/plant. Maize cobs were collected at harvest time (30 days) after the second spray of labeled ethion, and dried for preparation of

the cake and determination of radioactivity.

### Extraction of Maize Seeds

Seeds were crushed and extracted through soxhlet with n-hexane for 24 hrs. The cake after evaporation of hexane was further extracted with methanol for 6 h.

### Radioactivity Measurements

The ethion bound residue in cake used in the experiment was determined as follows: 100 mg of cake was digested, using [1 mL Solusol (tissue and gel solubilizer), 1 mL 30 % H<sub>2</sub>O<sub>2</sub> and 70 µL glacial acetic at 40–50°C]. The radioactivity was counted by Liquid Scintillation Counting (LSC) and it was 1.6 ppm.

### Preparation of Aqueous Extract Rosemary Leaves

Fresh rosemary leaves were collected from local market. The leaves were washed carefully, and then air dried in shade at room temperature, grinded to fine powder. Each 500 g of fine powdered herb was then refluxed in 2 liters of 70% ethanol at 60°-70°C for 36 hrs in a continuous extraction (soxhlet) apparatus. The ethanol extract was filtered and concentrated under reduced pressure at 60°C using a rotary evaporator. Ten grams of rosemary extract were dissolved in 100 mL of distilled water and incorporated in 1 kg of control and maize cake then mixed manually for at least 30 min. to ensure complete distribution.

### Toxicological Studies

Twenty male Wister rats, aged three months old and weighing 150 ± 20g, were obtained from the animal house of the National Research Centre, Dokki, Giza. Animals were maintained under controlled conditions of temperature at 25 ± 2°C, relative humidity of 50 ± 15%, and normal photoperiod (12h dark: 12h light) for one week giving them the standard rat pellet and water. They were provided with tap water and balanced diet ad libitum. All animals have received human care in compliance with the state authorities following the Egyptian rules of animal protection. The rats were randomly distributed into four groups of five rats each as follows:

Group 1: Animals fed on maize cake free from ethion residues or rosemary extracts (Control).

Group 2: Animals fed on maize cake free from ethion residues and rosemary extracts (1% w/w).

Group 3: Animals fed on maize cake containing ethion bound residues (1.6 ppm) (Treated).

Group 4: Animals fed on maize cake containing ethion bound residues (1.6 ppm) and rosemary extracts (1% w/w).

### Blood Sampling and Biochemical Analysis

At the end of the experimental period (30 days) the blood samples were collected from the eye plexuses of the animals. Then, it is divided into two parts, the first part was put into a clean centrifuge glass tube with EDTA to prepare plasma and the second part was put in a dry clean centrifuge glass tube without any coagulation to prepare serum. The samples were left for 15 minutes at normal of the temperature of room, after that the tubes were centrifuged for another 15 minutes at 3000 rpm for analysis of biochemical parameters. Cholinesterase activity was determined according to Ellman method [29] as modified by Gorun et al [30]. The liver parameters [alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP), total protein, and albumin], kidney parameters (urea and creatinine), lipid profile, [total cholesterol (TC), triglycerides (TG), high density lipoprotein (HDL), low density lipoprotein (LDL)] and antioxidant parameters [catalase (CAT) and glutathione-S-transferase (GST)] were determined using Kits purchased from Bio-diagnostic Co., Egypt.

### Histopathological Effect of Ethion on Selected Tissues

Selected tissues (testes, liver, kidney and brain) were isolated and fixed in 10% neutral buffered formalin for 24 hours, rinsed with water, dehydrated in alcohols, cleared in xylene, and embedded in paraffin. Tissue blocks were sectioned at 4-5 micron thickness and routinely stained with Haematoxylin and Eosin (H&E) stain [31] and examined by light microscopy.

### Statistical Analysis

The obtained data from serum biochemical and enzymes analysis were expressed as mean ± standard error. Data were analyzed by using Student's t-test. P < 0.05, P < 0.01 and P < 0.001 values were used [32].

### Results and Discussions

Ethion, an organophosphorus pesticide, used world widely causing public health concern [33]. Compared with control rats, the present results of the estimated biochemical parameters indicated that ethion bound residues caused a significant inhibition in the plasma acetylcholinesterase (P < 0.01) (Table 1). The inhibitory effect on acetylcholinesterase was expected, as the organophosphate compounds are anti cholinesterase agents for RBC, and plasma ChE, Such findings have been reported since the 1950's by many investigators [34,8,35]. Meanwhile, significant inhibition (P < 0.01) was continued when RM supplemented to ethion bound residues treated animals.

Groups	Plasma-Cholinesterase level ( $\mu$ mole/mL)
	Means a $\pm$ SD
Group 1	1.38 $\pm$ 0.040
Group 2	1.31 $\pm$ 0.055
Group 3	1.26 $\pm$ 0.034 ***
Group 4	1.242 $\pm$ 0.031***

**Table 1:** Effect of ethanolic extract of rosemary on cholinesterase activities of rats treated with ethion bound residues for 30 days. a: results are expressed as mean  $\pm$  SD for five samples, \*\*\*: Significance at  $P < 0.01$ .

The liver, as an organ of metabolism and excretion, is endowed with the task of detoxification. Toxicants such as viruses, fungal products, bacterial metabolites, pollutants and chemotherapeutic factors could cause different liver disorders [24]. Sensitive indicators of liver cell injury are serum transaminases such as AST, ALT and ALP. Our results have shown that oral administration of cake contaminated with ethion bound residues induced hepatotoxicity. The effects of subchronic administration of ethion bound residues and the protective effects of RM on serum biochemistry are shown in Table 2. Compared with control rats, activities of hepatic enzymes, ALT, AST, and ALP were significantly increased ( $P < 0.01$ ,  $P < 0.001$  and  $P < 0.01$ ), respectively (Group 3). Whereas the other two parameters, the total protein

and albumin were significantly decreased ( $P < 0.01$  and  $P < 0.05$ ), respectively. In contrast, RM-treatment significantly suppressed elevation of ALT, AST and ALP and increasing of the total protein and albumin concentrations (Group 4).

The liver is considered as one of the target organs affected by pesticides toxicity owing to its storage in the liver after pesticides exposure. In the current study, the liver enzymes assays indicated that ethion bound residues treatment induced significant elevation of serum AST, ALT, and ALP activities. It has been shown that organophosphorus insecticides can elevate the enzymatic activities of ALP, ALT, AST, and LDH [17,26]. Supplementation with RM leaves extract was able to diminish significantly the ethion-induced changes in all the mentioned parameters. RM leaves extract acts as an antioxidant with hepatoprotective effect which has been confirmed by Mehmetcik et al. These results are run in parallel with the results of [36] who found that, a significant improving effect of pre-treatment with rosemary on the altered activities of serum ALT, AST, GGT and ALP induced by Pb-acetate intoxication. Similarly, results of the study by Sotelo-Félix, et al. [20] and Gutiérrez, et al. [37] showed that methanolic extract of rosemary could restore the elevated ALT and AST activities in both acute liver damage and liver cirrhosis experimental models. Also, these results are in agreement with Albasha and Azab [38] who reported that aqueous extract of rosemary has significantly decreased the release of AST and ALT.

Groups	ALT (U/L)	AST (U/L)	(ALP) (U/L)	T. P (g/dL)	ALB (g/dL)
	Means a $\pm$ SD	Means a $\pm$ SD	Means a $\pm$ SD	Means a $\pm$ SD	Means a $\pm$ SD
Group 1	50 $\pm$ 8.09	110.4 $\pm$ 7.7	185.8 $\pm$ 15.5	4.25 $\pm$ 0.5	2.27 $\pm$ 0.46
Group 2	44.0 $\pm$ 6.16	108.6 $\pm$ 9.32	183.6 $\pm$ 8.88	4.27 $\pm$ 0.23	2.11 $\pm$ 0.15
Group 3	71.8 $\pm$ 5.26***	156.8 $\pm$ 8.35****	239.6 $\pm$ 5.73***	3.1 $\pm$ 0.1***	1.53 $\pm$ 0.1*
Group 4	51.0 $\pm$ 7.35***	104.8 $\pm$ 5.76****	199.2 $\pm$ 9.65***	4.17 $\pm$ 0.14****	2.06 $\pm$ 0.13***

**Table 2:** Effect of ethanolic extract of rosemary on liver function of rats treated with ethion bound residues for 30 days. a: results are expressed as mean  $\pm$  SD for five samples, \*: Significance at  $P < 0.05$ , \*\*\*: Significance at  $P < 0.01$ , \*\*\*\*: Significance at  $P < 0.001$ .

The elevated serum levels of urea and creatinine indicate reduced ability of the kidney to eliminate the toxic metabolic substances [39]. Many pesticides can cause some toxic and adverse effects on the kidney tissues [29]. Kidney is one of the target organs of experimental animals attacked by OP compounds and can alter plasma urea, uric acid, and creatinine levels [40]. The levels of urea and creatinine in the plasma of rats are measured as indicators for kidney function [6]. In this work, exposure to cake containing ethion bound residues (group 3) produced a significant increase ( $P < 0.01$ ) in the values of urea, and creatinine concentrations relative to the control group (Table 3) confirming a pronounced

impairment in kidney function.

Elevated serum creatinine level and blood urea were noticed in mice receiving ethion residues for three months [41]. These results agreed with the results of Abdel-Gawad, et al. [35] who illustrated that the exposure to ethion residues in soybean seeds increases the level of serum urea and creatinine. On the other hand, rats administered RM alone revealed insignificant change in the mentioned parameters as compared to the control group. The treatment with RM recorded significant ( $P < 0.001$ ) decrement in urea and creatinine in comparison with cake containing ethion bound

residues treated group (Table 3). These results suggest the ability of RM extract to restore impaired renal excretory function. This is in accordance with results of the study by Sakr and Lamfon [42], indicating that water extract of rosemary could reduce the levels of urea and creatinine in serum of rats, when compared to  $\text{CCl}_4$ -treated group of animals.

Group	Urea (mg %)	Creatinine (mg %)
	Means a $\pm$ SD	Means a $\pm$ SD
Group 1	59.0 $\pm$ 4.0	1.06 $\pm$ 0.64
Group 2	49.4 $\pm$ 5.73	1.002 $\pm$ 0.018
Group 3	76.8 $\pm$ 5.36***	1.55 $\pm$ 0.20***
Group 4	45.6 $\pm$ 4.77****	0.98 $\pm$ 0.038***

**Table 3:** Effect of ethanolic extract of rosemary on kidney function of rats treated with ethion bound residues for 30 days. a: results are expressed as mean  $\pm$  SD for five samples, \*\*\*: Significance at  $P < 0.01$ , \*\*\*\*: Significance at  $P < 0.001$ .

Groups	Cholesterol (mg %)	Triglycerides (mg %)	HDL (mg %)	LDL (mg %)
	Means a $\pm$ SD	Means a $\pm$ SD	Means a $\pm$ SD	Means a $\pm$ SD
Group 1	76.8 $\pm$ 5.07	71.0 $\pm$ 5.57	34.5 $\pm$ 1.73	14.0 $\pm$ 1.0
Group 2	71.6 $\pm$ 5.22	63.0 $\pm$ 2.45	40.2 $\pm$ 1.60	10.04 $\pm$ 0.31
Group 3	107.6 $\pm$ 7.58***	95.4 $\pm$ 5.55***	30.88 $\pm$ 1.6****	24.9 $\pm$ 1.67***
Group 4	72.2 $\pm$ 4.92****	61.6 $\pm$ 3.21****	36.66 $\pm$ 0.96****	15.26 $\pm$ 0.32****

**Table 4:** Effect of ethanolic extract of rosemary on lipid profile of rats treated with ethion bound residues for 30 days. a: results are expressed as mean  $\pm$  SD for five samples, \*\*\*: Significance at  $P < 0.01$ , \*\*\*\*: Significance at  $P < 0.001$ .

CAT and GST, constitute a mutually supportive team of defense against reactive oxygen species (ROS). CAT helps in the removal of  $\text{H}_2\text{O}_2$  formed during the reaction catalyzed by SOD. GST catalyzes the formation of the thiol group of glutathione to electrophilic xenobiotics. These antioxidant enzymes can, therefore, alleviate the toxic effects of ROS. In general, pesticide treatment produces oxidative stress by the generation of free radicals and induced tissue lipid peroxidation in mammals and other organisms. Reduced activities of antioxidant enzymes (CAT, SOD) after treatment of pesticides are important indicators for the toxicity of these chemicals [51-53]. The results obtained clearly indicated that treatment with ethion bound residues for 30 days resulted in a significant decrease in the activities of CAT and GST as compared to control animals (Table 5). However, rats treated with rosemary extract showed increase in CAT and GST as compared to control rats. When rosemary extract administered with ethion bound residues, it has capable of recovering the activities of CAT and GST to the normal values. Our results are in agreement with those found by Khalil, et

Considering that liver damage could result in a change of the lipid content in serum, cholesterol, triglycerides, HDL and LDL in serum were determined. Treatment with cake containing ethion bound residues (group 3) caused a significant ( $P < 0.01$ ) increase in cholesterol, triglyceride and LDL, while caused a significant decrease ( $P < 0.001$ ) in HDL content as compared to the control (Table 4). This is in agreement with recent studies on  $\text{CCl}_4$  poisoning [43]. RM extract supplementation also resulted in the significant attenuation in the level of serum, cholesterol, triglycerides, HDL and LDL in serum, toward the control level which again strengthens the hypolipidemic effect of this extract. Several studies have demonstrated a significant increase in the serum-lipid constituents in the experimental animals, treated with different OPI [44-46]. The hypolipidemic effects of RM in our studies were previously reported by Ozkol, et al. [47] and Labban, et al. [48], respectively. Other reported investigations are consistent with our results. Also, treatment rosemary oil in rats fed high fat diet modulated the elevation of lipids parameters [49-50].


al. [54] who reported that Serum enzymatic GST, (CAT), and (GPx) were found to be increased by the administration of *R. officinalis* in animals induced by Streptozotocin (STZ). Moreover, Abdel-Gawad, et al. [35] demonstrated that CAT and GST were found to be decreased in animals treated with  $^{14}\text{C}$ -Ethion residues and increased by the administration of artichoke leaf powder.

Groups	CAT (U/l)	GST(U/mL)
	Means a $\pm$ SD	Means a $\pm$ SD
Group 1	348 $\pm$ 15.03	3006 $\pm$ 99.52
Group 2	355 $\pm$ 15.17	3056 $\pm$ 70.88
Group 3	312 $\pm$ 6.57 ***	2523 $\pm$ 63.77****
Group 4	345 $\pm$ 11.51 ***	2922 $\pm$ 49.96 ****

**Table 5:** Effect of ethanolic extract of rosemary on Antioxidant activities of rats treated with ethion bound residues for 30 days. a: results are expressed as mean  $\pm$  SD for five samples, \*\*\*: Significance at  $P < 0.01$ , \*\*\*\*: Significance at  $P < 0.001$ .

## Histopathological Effect of Ethion on Selected Tissues


The histopathological examination of control liver sections showed normal histological structure (Figure 2 A,B). animals exposed to bound ethion showed mild congestion of central veins associated with mild mononuclear inflammatory cells infiltration (Figure 2C), which is apparently improved when using Rosemary (Figure 2 D) The histopathological examination of liver revealed severe congestion of central veins and hepatic sinusoids, associated with severe necrosis, vacuolar degeneration of hepatocytes and mononuclear inflammatory cells infiltration due to pesticide treatment as documented previously. The exposure to the pesticides impair the liver function and to cause injury to the hepatic tissues. The drastic alterations observed in the liver tissues due to pesticide exposure suggested hepatotoxic effects as reported by Iseri, et al. [55-57].




**Figure 2:** Photomicrographs of H and E stained Liver sections. (Figure A): Control rat shows normal histologic structure of liver (X 200). (Figure B): shows mild congested central veins (Arrows) (X 100). (Figure C): Shows mild congestion of central veins with mild inflammatory cells infiltration in sections exposed to Bound Ethion (X 100). (Figure D): shows apparent normal hepatic sections exposed to both Bound Ethion and Rose Mary extract. (X 200).

Control kidney sections showed normal histological features (Figure 3A). Examined sections of male rats treated with bound ethion showed mild hemorrhage within renal tissue (Figure 3B) as well as mild degenerative changes (Figure 3C) Which appeared as control ones (Figure 3D). The most prominent histopathological alterations of total ethion treated rats showed severe interstitial nephritis, hemorrhage and edema, associated with severe infiltration with inflammatory cells as previously concluded by Mamun et al [58-61]. Sections from control and Rosemary treated groups

showed no histopathological alterations in the cerebral cortex (Figure 4A) Treatment with bound ethion showed mild congestion of cerebral blood vessels associated with mild gliosis (Figure 4B, C). Brain of rat treated with bound ethion and Rosemary showing significant improvement with slight congestion of cerebral blood vessels (Figure 4D).



**Figure 3:** Photomicrographs of H and E stained Kidney sections. (Figure A): Control rat shows normal histologic structure of kidney. (Figure B): shows mild hemorrhage within renal tissue. (Figure C): shows mild degenerative changes in sections exposed to Bound Ethion. (Figure D): shows apparent normal kidney sections exposed to both Bound Ethion and Rosemary extract. (X 200).



**Figure 4:** Photomicrographs of H and E stained Brain sections. (Figure A): photomicrograph of control rat brain shows nohistopathologicstructure in cerebral cortex. (Figure B): shows mild congestion of cerebral blood vessels. (Figure C): Shows vacuolar degeneration in brain exposed to bound Ethion (Arrows). (Figure D): brain of rat treated with bound Ethion and Rosemary showing significant improvement with slight congestion of cerebral blood vessels (H& E X 100).

Ethion showed important degenerative changes in brain

tissue in the form of congestion of cerebral blood vessels, edema vacuolations as mentioned by Abdel-Salam, et al. [59] who stated that treatment with only OPIs causes' neuronal damage in the cerebral cortex and degeneration of some Purkinje cells in the cerebellum. The oxidative stress increased in the brain tissue suggesting increased generation of free radicals leading to neuronal injury [62-66] as well as decreased antioxidant enzyme activities [67-69]. The administration of Rosemary has been reported to attenuate and ameliorate the alterations caused by ethion in male rat.

## Conclusion

In conclusion, the present data revealed that RM extract had free radical scavenging capacity. Ethion bound residues administration induced liver and kidney damage as well as hyperlipidemic effects as evidenced from elevated hepatic, kidney and lipidemic markers. However, coadministration of RM extract markedly ameliorated the deleterious effects of ethion bound residues mainly through their antioxidant activity. Histopathological studies on liver, kidney and brain tissues confirmed our data. It is possible that RM extract may be useful as adjuvant due to reduce some of the adverse effects of the pesticide bound residues. Thus, this study suggests the potent role of rosemary in management of injury-induced by ethion bound residues exposure and this effect may be attributed to its antioxidant activity.

## References

- Alexandratos Nikos, Jelle Bruinsma (2012) World agriculture towards 2030/2050 the revision.
- Damalas CA, Eleftherohorinos IG (2011) Pesticide exposure, safety issues, and risk assessment indicators. *Int J Environ Res public Health* 8(5): 1402-1419.
- John EC, Kathleen A Durkin (2013) Anticholinesterase insecticide retrospective. *Chemico-biological interactions* 203(1): 221-225.
- Soderlund DM (2012) Molecular mechanisms of pyrethroid insecticide neurotoxicity: recent advances. *Arch Toxicol* 86(2): 165-181.
- Gee David, Philippe Grandjean, Steffen Foss Hansen, Malcolm MacGarvin, Jock Martin, Gitte Nielsen, et al. (2013). Late lessons from early warnings: science, precaution, innovation. pp: 746
- Kapoor SK, Kalra RL (1997) Transfer of HCH Isomers from Feed into Milk of Indian Buffalo *Bubalus hubalis L* *Pesti Res J* 9(1): 72-78.
- Tunçbilek AŞ, P Aysal, and MB Halitligil (1997) Loss of label during processing cotton oil grown with 14 C-aldicarb. *Bull Environ Contam Toxicol* 58(2): 213-218.
- Bhatti GK, Kiran R, Sandhir R (2010) Modulation of ethion-induced hepatotoxicity and oxidative stress by vitamin E supplementation in male Wistar rats. *Pesticide biochemistry and physiology* 98(1): 26-32.
- Howard PhilipH (2017) Handbook of environmental fate and exposure data: for organic chemicals, volume III pesticides. Routledge.
- Desouky MM, Abdel-Gawad H, Hegazi B (2013) Distribution, fate and histopathological effects of ethion insecticide on selected organs of the crayfish, *Procambarus clarkii*. *Food chem toxicol* 52: 42-52.
- Hosseinimehr SJ (2014) Beneficial effects of natural products on cells during ionizing radiation. *Rev Environ Health* 29(4): 341-353.
- Ibarra A, Cases J, Bily A, He K, Bai N, et al. (2010) Importance of extract standardization and in vitro/ex vivo assay selection for the evaluation of antioxidant activity of botanicals: a case study on three *Rosmarinus officinalis L.* extracts. *J med food* 13(5): 1167-1675.
- Erkan N, Ayranci G, Ayranci E (2008) Antioxidant activities of rosemary (*Rosmarinus Officinalis L.*) extract, blackseed (*Nigella sativa L.*) essential oil, carnosic acid, rosmarinic acid and sesamol. *Food chem* 110(1): 76-82.
- Nolkemper S, Reichling J, Stintzing FC, Carle R, Schnitzler P (2006) Antiviral effect of aqueous extracts from species of the Lamiaceae family against Herpes simplex virus type 1 and type 2 in vitro. *Planta Med* 72(15): 1378-1382.
- Del Bano MJ, Castillo J, Benavente-García J, Lorente J, Martín-Gil R, et al. (2006) Radioprotective-Antimutagenic Effects of Rosemary Phenolics against Chromosomal Damage Induced in Human Lymphocytes by  $\gamma$ -rays. *J Agric Food Chem* 54(6): 2064-2068.
- Bernardes WA, Lucarini R, Tozatti MG, Flauzino LG, Souza MG, et al. (2010) Antibacterial Activity of the Essential Oil from *Rosmarinus offi cinalis* and its Major Components against Oral Pathogens. *Z Naturforsch C J Biosci* 65(9-10): 588-593.
- Chen CC, Chen HL, Hsieh CW, Yang YL, Wung BS (2011) Upregulation of NF-E2-related factor-2-dependent glutathione by carnosol provokes a cytoprotective response and enhances cell survival. *Acta Pharmacol Sin* 32(1): 62.
- Berrington D, Lall N (2012) Anticancer activity of certain

- herbs and spices on the cervical epithelial carcinoma (HeLa) cell line. Evid Based Complement Alternat Med.
19. Güllüce M, Sökmen M, Daferera D, Açar G, Özkan H, et al. (2003) In vitro antibacterial, antifungal, and antioxidant activities of the essential oil and methanol extracts of herbal parts and callus cultures of *Satureja hortensis* L. J Agric Food Chem 51(14): 3958-3965.
  20. Takaki I, LE Bersani-Amado, A Vendruscolo, SM Sartoretto, SP Diniz, et al. (2008) Anti-inflammatory and antinociceptive effects of *Rosmarinus officinalis* L. essential oil in experimental animal models. J Med Food 11(4): 741-746.
  21. Afonso MS, de O Silva AM, Carvalho EB, Rivelli DP, Barros SB, et al. (2013) Phenolic compounds from Rosemary (*Rosmarinus officinalis* L.) attenuate oxidative stress and reduce blood cholesterol concentrations in diet-induced hypercholesterolemic rats. Nutr metab 10(1): 19.
  22. Ramadan KS, Khalil OA, Danial EN, Alnahdi HS, Ayaz NO (2013) Hypoglycemic and hepatoprotective activity of *Rosmarinus officinalis* extract in diabetic rats. J Physiol Biochem 69(4): 779-783.
  23. Azab AE, Fathy Ahmed Fetouh, Mohamed OA (2014) Nephro-protective effects of curcumin, rosemary and propolis against gentamicin induced toxicity in guinea pigs: Morphological and biochemical study. Ameri J Clin Experi Medi 2(2): 28-35.
  24. Haraguchi H, Saito T, Okamura N, Yagi A (1995) Inhibition of lipid peroxidation and superoxide generation by diterpenoids from *Rosmarinus officinalis*. Planta Med 61(4): 333-336.
  25. Amin A, AA Hamza (2005) Hepatoprotective effects of Hibiscus, *Rosmarinus* and *Salvia* on azathioprine-induced toxicity in rats. Life sci 77(3): 266-278.
  26. Munné-Bosch S, Schwarz K, Alegre L (1999) Enhanced formation of  $\alpha$ -tocopherol and highly oxidized abietane diterpenes in water-stressed rosemary plants. Plant Physiol 121(3): 1047-1052.
  27. Gallego M Gabriela, Michael H Gordon, Francisco J Segovia, Monika Skowrya, María P Almajano (2013) Antioxidant properties of three aromatic herbs (rosemary, thyme and lavender) in oil-in-water emulsions. J Americ Oil Chem Society 90(10): 1559-1568.
  28. Abdel-Gawad H, RM Abdelhameed, AM Elmesalamy, B Hegazi (2011) Distribution and elimination of  $^{14}\text{C}$ -ethion insecticide in chamomile flowers and oil. Phosphorus, Sulfur, and Silicon and the Related Elements 186(10): 2122-2134.
  29. Ellman, George L, K Diane Courtney, Valentino Andres Jr, Robert M Featherstone (1961) A new and rapid colorimetric determination of acetylcholinesterase activity. Biochem pharmacol 7(2): 88-95.
  30. Gorun V, Proinov I, Băltescu V, Balaban G, Bârzu O (1978) Modified Ellman procedure for assay of cholinesterases in crude enzymatic preparations. Anal Biochem 86(1): 324-326.
  31. Bancroft, John D, Marilyn Gamble (2008) Theory and practice of histological techniques: Elsevier health sciences.
  32. Snedecor, George W, and WG Cochran (1980) Statistical methods, 7<sup>th</sup> (Ed.), Ames, Iowa State University Press, USA.
  33. Verma G, Chandra SM, Ramneek V, Baljit S, Sethi RS (2019) Long-term exposures to ethion and endotoxin cause lung inflammation and induce genotoxicity in mice. Cell and tissue research 375(2): 493-505.
  34. Nigg HN, Knaak JB (2000) Blood cholinesterases as human biomarkers of organophosphorus pesticide exposure. Rev Environ Contam Toxicol 163: 29-111.
  35. Abdel-Gawad Hassan, Hamdy Taha, Reda Abdel-Hameed, Bahira Hegazi (2013)  $^{14}\text{C}$ -Ethion residues in soybean seeds: metabolic pathway, effect of processing, bioavailability, toxicity and protective action of artichoke leaf powder towards rats. Toxil Envi Chem 95(2): 288-303.
  36. Abd El Kader MA, NM El-Sammad, H Taha (2012) The protective role of rosemary (*Rosmarinus officinalis*) in lead acetate induced toxicity in rats. J Appl Sci Res 8(6): 3071-3082.
  37. Ha KT, Yoon SJ, Choi DY, Kim DW, Kim JK, et al. (2005) Protective effect of *Lycium chinense* fruit on carbon tetrachloride-induced hepatotoxicity. J Ethnopharmacol 96(3): 529-535.
  38. Albasha Mohamed Omer, SA Azab (2014) Effect of cadmium on the liver and amelioration by aqueous extracts of fenugreek seeds, rosemary, and cinnamon in Guinea pigs: histological and biochemical study. Cell Biol 2(2): 7-17.
  39. Mehmetçik G, Ozdemirler G, Koçak-Toker N, Cevikbaş U, Uysal M (2008) Effect of pretreatment with artichoke extract on carbon tetrachloride-induced liver injury and oxidative stress. Exp Toxicol pathol 60(6): 475-480.



40. EL, SABAH G, AHMED M, ATTIA, AFAF M (2009). Effect of garlic consumption on blood lipid and oxidant/antioxidant parameters in rat males exposed to chlorpyrifos. *Slovak J Anim Sci* 42(3): 111-117.
41. Sakr SA, Lamfon HA (2012) Protective effect of rosemary (*Rosmarinus officinalis*) leaves extract on carbon tetrachloride-induced nephrotoxicity in albino rats. *Life Sci J* 9(1): 779-785.
42. El-Hadary, Abdalla E, Elsanhoty RM, Ramadan MF (2019) In vivo protective effect of *Rosmarinus officinalis* oil against carbon tetrachloride (CCl<sub>4</sub>)-induced hepatotoxicity in rats. *PharmaNutrition* 9: 100151.
43. Ibrahim NA, El-Gamal BA (2003) Effect of diazinon, an organophosphate insecticide, on plasma lipid constituents in experimental animals. *J Biochem Mol Biol* 36(5): 499-504.
44. Lasram MM, Annabi AB, El Elj N, Selmi S, Kamoun A, et al. (2009) Metabolic disorders of acute exposure to malathion in adult Wistar rats. *J Hazard Mater* 163(2-3): 1052-1055.
45. Yehia, Mona AH, Sabah G El-Banna, Aly B Okab (2007) Diazinon toxicity affects histophysiological and biochemical parameters in rabbits. *Exper Toxicol pathol* 59(3-4): 215-225.
46. Ozkol H, Tuluçe Y, Dilsiz N, Koyuncu I (2013) Therapeutic potential of some plant extracts used in Turkish traditional medicine on streptozocin-induced type 1 diabetes mellitus in rats. *J Membr Biol* 246(1): 47-55.
47. Labban L, El-Sayed MU, Yasser MI (2014) The effects of rosemary (*Rosmarinus officinalis*) leaves powder on glucose level, lipid profile and lipid peroxidation. *Inter J Clini Med* 5(06): 297-304.
48. Nazni PT, Vijayakumar P, Alagianambi P, Amirthaveni M (2006) Hypoglycemic and Hypolipidemic Effect of *Cynara Scolymus* among Selected Type 2 Diabetic Individuals. *Pak J Nutr* 5(2): 147-151.
49. Eissa FA, Choudhry H, Abdulaal WH, Baothman OA, Zeyadi M, et al. (2017) Possible hypocholesterolemic effect of ginger and rosemary oils in rats. *Afr J Tradit Complement Altern Med* 14(4): 188-200.
50. Altuntas I, Delibas N, Demirci M, Kilinc I, Tamer N (2002) The effects of methidathion on lipid peroxidation and some liver enzymes: role of vitamins E and C. *Arch Toxicol* 76(8): 470-473.
51. Hazarika A, Sarkar SN, Hajare S, Kataria M, Malik JK (2003) Influence of malathion pretreatment on the toxicity of anilofos in male rats: a biochemical interaction study. *Toxicology* 185(1-2): 1-8.
52. Jayakumar T, Ramesh E, Geraldine P (2006) Antioxidant activity of the oyster mushroom, *Pleurotus ostreatus*, on CCl<sub>4</sub>-induced liver injury in rats. *Food and chem Toxicol* 44(12): 1989-1996.
53. Olfat AK, Kholoud SR, Enas ND, Hanan SA, Najla OA (2012) Antidiabetic activity of *Rosmarinus officinalis* and its relationship with the antioxidant property. *Afri J Phar Pharmacol* 6(14): 1031-1036.
54. Vivek GK, Nikhat JS, Ojha AK, Bechan S (2019) Hepatoprotective effect of Aloe vera against cartap-and malathion-induced toxicity in Wistar rats. *J cellul physiol* 234(10): 18329-18343.
55. El-Beshbishy, HA (2005). Hepatoprotective effect of green tea (*Camellia sinensis*) extract against tamoxifen-induced liver injury in rats. *BMB Reports* 38(5): 563-570.
56. Işeri S, Ercan F, Gedik N, Yüksel M, Alican I (2007) Simvastatin attenuates cisplatin-induced kidney and liver damage in rats. *Toxicology* 230(2-3): 256-264.
57. Mamun AA, Rahman A, Belal SH, Islam MA, Sarker MEH, et al. (2015) Histological Study of the Effect of Malathion on Liver and Kidney Tissues of Mice Model. *Inter J Pharmace Sci Res* 6(3): 1043-1048.
58. Abdel-Salam Omar ME, Amany A Sleem, Eman R Youness, Fatma A Morsy (2018) Preventive effects of cannabis on neurotoxic and hepatotoxic activities of malathion in rat. *Asian Pacific J Trop Med* 11(4): 272-279.
59. EI\_Bini Dhouib, Lasram MM, Annabi A, Gharbi N, El-Fazaa S (2015) A comparative study on toxicity induced by carbosulfan and malathion in Wistar rat liver and spleen. *Pestic Biochem Physiol*.
60. Karabag-Coban F, Ibrahim B, Liman R, Sinan I, Cigerci I, et al. (2016) Oleuropein alleviates malathion-induced oxidative stress and DNA damage in rats. *Toxicol Environ Chem* 98(1): 101-108.
61. Abdel-Salam Omar ME, Eman R Youness, Nadia A Mohammed, Noha N Yassen, Yasser A Khadrawy, et al. (2016) Novel neuroprotective and hepatoprotective effects of citric acid in acute malathion intoxication. *Asian Pac J Trop Med* 9(12): 1181-1194.
62. Abdel-Salam OME, Youness ER, Mohammed NA, Yassen NN, Khadrawy YA, et al. (2017) Nitric oxide synthase inhibitors protect against brain and liver damage caused by acute malathion intoxication." *Asian Pac J Trop Med* 10(8): 773-786.

63. Celik I, Yilmaz Z, Turkoglu V (2009) Hematotoxic and hepatotoxic effects of dichlorvos at sublethal dosages in rats. *Environ Toxicol*: 24(2): 128-132.
64. Elhalwagy Manal EA, Nevine S Darwish, Enass M Zaher (2008) Prophylactic effect of green tea polyphenols against liver and kidney injury induced by fenitrothion insecticide. *Pesticide biochem physiology* 91(2): 81-89.
65. Gutiérrez R, Alvarado JL, Presno M, Pérez-Veyna O, Serrano CJ, et al. (2010) Oxidative stress modulation by *Rosmarinus officinalis* in CCl<sub>4</sub>-induced liver cirrhosis. *Phytother Res* 24(4): 595-601.
66. Hummadi Laila A (2012) Histopathological and ultrastructural changes in renal corpuscle of female rats topical application by p-phenylene diamine. *Inter J Zoologi Res* 8(3): 106-120.
67. Kaneko J, John H, Michael B (2008) Clinical biochemistry of domestic animals. Academic press.
68. Sameeh MA, Abdel-Tawab HM (2010) Oxidative damage, biochemical and histopathological alterations in rats exposed to chlorpyrifos and the antioxidant role of zinc. *Pest Biochem Physiol* 96(1): 14-23.
69. Sotelo-Felix JI, D Martinez-Fong, P Muriel, RL Santillan, D Castillo, et al. (2002) Evaluation of the effectiveness of *Rosmarinus officinalis* (Lamiaceae) in the alleviation of carbon tetrachloride-induced acute hepatotoxicity in the rat. *J Ethnopharmacol* 81(2): 145-154.