



Studies on the Definition of Rodent Species at Farshut District, Qena Governorate, Egypt

Elrawy AAA¹, Mahmoud NA¹, Baghdadi SAS¹ and Abd El-Aleem SSD^{2*}

¹Agric. Zoology and Nematology Dept, Al-Azhar University, Egypt

²Plant Protection Department, Faculty of Agriculture Sohag University, Egypt

***Corresponding author:** Abd El-Aleem SS Desoky, Plant protection Department, Faculty of Agriculture, Sohag University, Egypt, Email: abdelaleem2011@gmail.com

Received Date: March 12, 2021; **Published Date:** March 26, 2021

Abstract

This study was carried out to identify rodent species in the old areas of Al-Dahsa village in Farshout district, Qena governorate, in field crops and buildings during two consecutive years (2018 and 2019). The study revealed the presence of two rodent species (such as *Arvicanthis niloticus* and *Rattus rattus frugivorus*) in the field crops. While the study revealed the presence of three species of rodents (*Mus musculus*, *Rattus rattus frugivorus*, and *Rattus norvegicus*) in the buildings. The study also revealed an increase in the density of rodents in field crops in the first and second years as follow: *R. r. frugivorus* (55.46 and 55.65%), *A. niloticus* (44.54 and 44.35%) for the first and second year, respectively. In buildings, the domestic mouse *M. musculus* was the most widespread species, accounting for 48.39% and 44.44% of the total rodents trapped for each of the first and second years. It is followed by *Rattus norvegicus* with 29.03% and 34.19%. *R. r. frugivorus* was the last species occurring 23.39% and 21.37%. This study is useful for choosing the most appropriate control methods for the rodent species spread in the study area.

Keywords: Field Crops; Buildings; *Arvicanthis Niloticus*; *Rattus R Frugivorus*; *Rattus Norvegicus*; *M Musculus*

Introduction

Rodents are global in their distribution and they constitute the largest group of mammals represent nearly 40%, of all mammals living at the present time Bajomi [1]. Hoogstral, et al., [2] surveyed 51 species of rodent in Egypt, belonged to sub-order; Myomorpha, eleven species fall under family Muridae, subfamily; Murinae (genera: *Arvicanthis*, *Rattus*, *Acomys*, *Mus* and *Nesoke*) are domestic and commensal animals found abundance, while five families are low abundance in desert and semi-desert. Many researchers found in Upper Egypt, Beni-Suif, El Minia, Assiut, Sohag and Qena Governorates the dominant rate species were, *R. norvegicus*, *R. rattus*, *A. cahirinus*, *A. niloticus*, *M. musculus*, *Geribilus spp.* and *Juculus spp.* The dominant species and density related to habitats, crop installation, nearly reclamation land and abundance shelter and food and

seasons, Salit, et al., [3], Abazid [4], Hussein [5]; El-Deep and Lokma [6]; Embarak [7] and Desoky 2007 [8]. The aim of the current study is to know the rodents scattered in the center of Farshout, Qena Governorate.

Materials and Methods

Survey of rodents in old areas at EL-Dahasa village (located at 76 km. North of Qena Governorate), Farshut district, Qena Governorate was carried out in the field crops and a building was carried out during two successive years (2018 and 2019). This area has been planted from a long period with isolated patches of vegetables, wheat, Egyptian clover, palm trees, cattle farm, alfalfa and different orchards as well as a building and a poultry farm. The total wire-box traps were 50 wire-box traps of the usual spring door (25×12×10 cm) in the month. Were baited and distributed at 6pm and

examined each morning and the bait was replaced by fresh one. Trapped rodents were transferred to the laboratory for identified according to Osborn and Helmy [9].

Results and discussion

The study revealed the occurrence of two rodent species in field crops (viz., *Arvicanthis Nilotic us* and *Rattus rattus frugivorus*) these rodents belong to family Muridae, Suborder Myomorpha, order Rodentia Table 1 and Figure 1. This viz., similar with Elrawy [10] in sugar cane at Qena Governorate where found that the presence of species of rats included the white bellied rat, *R.r.frugivorus*, Nile grass rat, *A.niloticus* and *R.r.frugivorus* the dominant species, this may be due to the presence of attributed to the availability of food and shelter as well as prefers trees for nesting in houses. Also this may be due to the inter-specific competition between this species and other species. The study revealed the occurrence of three rodent species in buildings (viz., *Mus musculus*, *Rattus rattus frugivorus* and *Rattus norvegicus*) these rodents belong to family Muridae, Suborder Myomorpha, order Rodentia Table 1 and Figure 2. This data agree with that obtained by El-Khayat [11] in buildings at Qalubia Governorate (Tukh

district) where found that the presence of species of rats included *Mus musculus* (The house mouse), *Rattus rattus* (The climb rat) and *Rattus norvegicus* (The Norway rat). The abundance of rodent species in field crops in the first and second year. Could be arranged quantitatively in the following descending order:

- The White bellied rat, *R. r. frugivorus* (55.46 and 55.65% for the first and second year, respectively).
- The Nile grass rat, *A. niloticus* (44.54 and 44.35% for the first and second year, respectively).

In the buildings, the house mouse *M. musculus* was the most dominant species constituting 48.39% and 44.44% of the total trapped rodents for both first and second year. Followed by the Norway rat *Rattus norvegicus* 29.03% and 34.19%. The white bellied rat *R. r. frugivorus* was the last species of occurrence 23.39% and 21.37% (Table 1). It may be concluded that the availability of preferred food in both areas led the rats to feed, and produce higher reproduction rat in both areas, wholly in agreement with the findings obtained by Abdel-Gawad et al., [12] & El-Feky [13].

Area	Study of year	Species/ total	Arvicanthis niloticus		Rattus rattus frugivorus		Rattusnorvegicus		Mus musculus	
			No.	%	No.	%	No.	%	No.	%
Field crops	2018	119	53	44.54	66	55.46	0	0	0	0
	2019	115	51	44.35	64	55.65	0	0	0	0
Buildings	2018	124	0	0	29	23.39	36	29.03	59	47.58
	2019	117	0	0	25	21.37	40	34.19	52	44.44
Total		475	104	21.89	184	38.74	76	16	111	23.37

Table 1: Survey of rodent species captured from study areas during two successive years at Qena Governorate, during 2018-2019 years.




Figure 1: Percentage of rodent species in the field crops during two successive years at Qena Governorate.




Figure 2: Percentage of rodent species in the buildings during two successive years at Qena Governorate.

References

- Bajomi D (1984) Commensal rodent problems in Hungary Organization and practice of vertebrate pest control Hampshire Conference England. pp: 18-20.
- Hoogstral H (1963) A brief review of contemporary land mammals of Egypt (including Sinai). 2: lagomorpha and Rodentia. Journal of the Egyptian Public Health Association 38(1): 1-35.
- Salit AM, Helal TY, Ali MA, Abdel – Gawad KH, Arafa MS (1982) Composition of the rodent species in newly reclaimed semi-desert area comparing with cultivated land. Assiut Journal of Agric. Sciences 13(2): 53-62.
- Abazaid AA (1990) Efficiency of some common used rodenticide and some new alternatives against rodents in Qena Govern-orate. M.Sc. Thesis, Fac. Agric, Assiut Univ, pp: 93.
- Hussien SSM (1991) Ecological studies and control of certain rodents in Beni suef Governorate. M. Sc. Thesis, Fac. of Agric, Cairo Univ.
- El-Deeb HI, Lokma HE (1992) Field studies on population dynamics and reproductive biology of the Nile rat *Arvicanthis niloticus*. Food and Agriculture Organization of the United Nations 19(3): 1431-1435.
- Embarak MZ (1997) Ecological, control studies on Rodents and their ectoparasites in cultivated and Newly Reclaimed area. M.Sc. Thesis, Fac. Agric, Assiut Univ, pp: 130.
- Desoky ASS (2007) Management strategies for rodents within different Ecosystems. M.Sc. Thesis, Fac Agric Assiut Univ, Assiut, Egypt. pp: 124.
- Osborn DJ, Helmy I (1980) The contemporary land mammals of Egypt (including Sinai). Field Museum of Natural History. Fieldiana. Zoology; new ser, no. 5, pp: 569.
- Elrawy AAA (2017) Ecological Studies on some rodents caused damage on sugar crops at Assiut and Qena Governorates and its control. M.sc. Fac Agric Al-Azhar Univ pp: 112.
- El-Khayat AEE (2010) Ecological and biological studies on some species of rodents. M. Sc. Thesis, Fac. Agric, Benha Univ, Benha, Egypt pp: 161.
- Abdel-Gawad KH, Farghal AI, Salman AG, Ali AM (1987) Survey and population density of rodent species in some urban, rural and cultivated areas in Sohag Governorate, Upper Egypt. Assiut J Agric Sci 18(1): 259-266.
- El-Feky MA (1990) Studies on small rodents and their fleas species of public health importance in Sabahiya Experimental Station with special reference to their control. M.Sc. Thesis, Fac Agric Ale Univ, Egypt. pp: 113.