



Role of Some Phytohormones in Plant Response to Salt Stress

Abobatta WF*, Nadia AM and Hamed

Department of Citrus Research, Horticulture Research Institute, Egypt

***Corresponding author:** Waleed Fouad Abobatta, Department of Citrus Research, Horticulture Research Institute, Agriculture Research Center, Giza, Egypt, Tel: +201224296948; Email: wabobatta@arc.sci.eg

Received Date: May 07, 2024; **Published Date:** June 24, 2024

Abstract

Plants are extremely subjected to environmental stress, which includes drought, low or high temperatures, and salt stress. The most common environmental stress factor is salinity, primarily in arid and semi-arid regions. Salinity stress is the most limiting factor for seed germination and plant productivity; therefore, most plants are susceptible to adverse effects because of the high concentration of salt in the soil, which affects plant growth and productivity. Phytohormone, considered an endogenous defense factor under stress conditions, has an important role in plant growth and survival, particularly under abiotic stress. Phytohormones influence a wide spectrum of processes in plants in both normal and stressful environments. The phytohormones stimulate plant responses to stress conditions, such as jasmonic acid, while there are other phytohormones that include Proline, Gibberellic acid, Cytokinins and Auxins. Exogenous application of plant growth regulators could be an alternate strategy to improve salt stress tolerance. The use of auxin or gibberellin has been shown to mitigate the violent effects of abiotic stresses on seed germination, plant growth, and productivity. Furthermore, proline induces plant tolerance against various abiotic stresses by increasing their endogenous levels and their intermediate enzymes in plants. Moreover, exogenous application of proline may be a good approach to reducing the unfavorable effects of salinity stress on plants.

Keywords: Auxin; Cytokinins; Gibberellin; Phytohormones; Proline; Salinity

Abbreviations: ACC: 1-Amino Cyclopropane 1-Carboxylic Acid; ACO: ACC Oxidase; ACS: ACC Synthase; ATPS: ATP-Sulfurylase; AVG: Amino Ethoxy Vinyl Glycine; BR: Brassinosteroids; Ca: Calcium; CAT: Catalase; CBL: Calcineurin B-Like Protein; CK: Cytokinins; EBL: Epibrassinolide; GA: Gibberellins; GS: Glutamine Synthase; GSA: Glutamate-Semialdehyde; IAA: Indole Acetic Acid; JA: Jasmonate; MDA: Malondialdehyde; MeJA: Methyl Jasmonate; N: Nitrogen; NiR: Nitrite Reductase; NO: Nitric Oxide; NR: Nitrate Reductase; NRA: Nitrate Reductase Activity; OAT: Ornithine-Delta-Aminotransferase; P5C: Pyrroline-5-Carboxylate; P5CDH: P5C Dehydrogenase; P5CR: P5C

Reductase; P5CS: 1-Pyrroline-5-Carboxylate Synthetase; PDH: Proline Dehydrogenase; POD: Peroxidase; POX: Proline Oxidase; ROS: Reactive Oxygen Species; S: Sulfur; SNP: Sodium Nitroprusside; SOD: Superoxide Dismutase; SOS: Salt Overly Sensitive.

Introduction

Climate change affects soil salinity, particularly in arid and semi-arid regions, due to low precipitation, which has negative impacts on plant growth and development. Fluctuations in climate conditions increase changes in the

composition and structure of various metabolite processes. Proteins and RNA molecules that precede signal transduction or stress acclimation events in plants [1]. Furthermore, abiotic stresses affect plant metabolism, physiology, and morphology processes and cause disturbances in various cellular pathways, particularly genetic regulation. One of these significant changes in environmental conditions is salinity. There is about 20% of the cultivated area, and nearly half of the world's irrigated lands are affected by salinity [2]. Salinity is considered one of the main threats that affect plant growth, seedling development, inhibiting seed germination, inhibiting enzyme activity, synthesis of DNA, RNA, and protein, and mitosis, as well as minimizing biomass accumulation [3,4].

In high salinity conditions, plants either decrease their growth and developmental programs in response to stress due to the effects of specific ions on metabolism or reverse water relations [5]. Therefore, under rising salinity, plants use different strategies to adapt to stress conditions, reducing adverse impacts on plant development, in addition to stress conditions pushing plants to regulate growth and development to complete their life cycle [3]. Phytohormones (both endogenous and exogenous applications) mediated the plant response and played an important role in acclimating plants to both biotic and abiotic unfavorable conditions [6].

Plant growth regulators play a paramount role in the growth, flowering, and fruit set of different crops, especially gibberellic acid and naphthalene acetic acid, which promote fruit set and decrease fruit drop in many fruit species and varieties under stress conditions [7].

Biochemical mechanisms are considered one of the major adaptation techniques. Plant phytohormones are involved as essential components in various metabolic processes in the plant. While the interaction between different phytohormones causes different results, it also plays an important role in plant responses to different abiotic stresses.

The main phytohormones that stimulate plant responses to various stresses are jasmonic acid (JA), salicylic acid (SA), and abscisic acid (ABA), while there are other phytohormones that include triazole (TR), proline, gibberellic acid (GA), cytokinins (CKs), Auxins, Indole-3-acetic acid (IAA), abscisic acid (ABA), and ethylene (ETHY) [8].

Phytohormones play essential roles in enhancing plant adaptation to harsh environmental conditions, such as soil salinity, drought, and rising temperature. They also stimulate the plant to grow under stress. There is crosstalk (antagonisms or associations) amongst different phytohormones depending on the growth stage, severity, and duration of stress conditions [9]. In addition, it has an

effective role in controlling growth and nutrient uptake to reduce the negative impacts of salinity on plant growth and complete its life cycle under stress conditions [10].

It is generally clear that the repressive effect of salinity on seed germination and plant growth could be related to a decline in endogenous levels of phytohormones. It is suggested that phytohormones play an important role in stress responses and adaptation, thus, some researchers have used plant growth regulators to reduce the adverse effects of salinity stress [11].

However, plants should manufacture low-molecular-weight non-enzymatic antioxidants like proline to save cells and tissues from oxidative injury, as well as enzymatic antioxidants including peroxidase, superoxide dismutase, ascorbate peroxidase, and catalase to safeguard against oxidative stress [12].

Objectives of this Review

The present work focuses on increasing our understanding of the role of phytohormones in salinity tolerance and improving plant growth and development under salinity stress.

Through discussion of the influence of salinity stress on plant growth and crop productivity, explore the role of phytohormones in reducing the harmful effects of salinity on plant growth and reproduction. In addition, this work further highlights the importance of using phytohormones to develop salinity-tolerant crops in light of the future impact of predicted climate change through an exploration of 200 references and a discussion of 90 references related to the main subject.

Salinity

In arid and semi-arid regions, salinity is one of the most significant abiotic threats influencing plant development and output. Salt problems are more severe in hot, dry weather than in cold, damp weather; moreover, salinity conditions reduce the formation of dry matter that is required for various growth processes [13].

Regularly, the rates of photosynthesis are lower in plants exposed to salinity due to the negative effect of salinity on photosynthesis, this would furthermore lead to restrictions on water availability and an imbalance in nutrient uptake by plants with inhibition in seed germination due to ionic disturbance, osmotic, and toxic effects [14]. While salt-tolerant plants (halophytes) involved fewer Na^+ and Cl^- transport to leaves, were able to preserve lower Na^+/K^+ proportions in their shoots, and developed higher facilities

to isolate ions into the cytoplasm or cell wall [15]. However, plant species vary in their sensitivity or tolerance to salinity stress, most of the economic plants are glycophytes, which are sensitive to salt stress. For instance, citrus is generally classified as a salt-sensitive crop because physiological trouble and decreasing growth and fruit yield can occur at relatively low-salinity levels [16].

Both osmotic stress and ionic toxicity are due to salt stress. Salt stress reduces osmotic potential which leads to osmotic stress, which limits the absorption of water from the soil, while ionic stress is caused by the over-accumulation of toxic salt ions within plant cells. Both abiotic stresses and nutrient deficiency affect the various physiological statuses and are responsible for adverse effects on the growth and development of glycophytes crops [17].

Soil salinity affects plant growth through reducing the osmotic potential of the soil solution, interfering with nutrient uptake, inducing ionic toxicity, and associating nutrient imbalances [18]. There are major variations in the morphology and anatomy of plants grown on saline soils, rising salinity reduces the economic productivity of various crops by inhibiting the germination of seeds, decreasing seedling flushing, and fluctuating flowering and fruit set, which affect total yield and fruit quality [19].

High salinity forces ionic and osmotic stresses on plants. In earlier stages of plant responses, a rapid osmotic phase is caused by the osmotic pressure of high salt concentrations in the soil. After salt accumulates above a threshold level in the plant, a slower ionic phase occurs because of the toxic effects of salt ions [20].

The decrease in leaf area, stomatal conductance, and chlorophyll levels in the plants and the generation of reactive oxygen species (ROS) in all these aspects are due to salinity stress [21].

The reduction of photosynthetic ability under salinity is due to stomata closure, loss of chlorophyll, and the deactivation of specific enzymes, which are involved in the synthesis of photosynthetic pigments. The decrease in chlorophyll contents is mainly due to the demolition of chlorophyll biosynthesis, chlorine accumulation, and reductions in magnesium, iron, and manganese [22]. Furthermore, increasing salt concentrations in the soil reduce plant water absorption capacity, hurt metabolic processes, and affect negatively osmotic balance, nutrient absorbance, hydraulic conductivity, net photosynthetic rate, and intercellular CO₂ concentrations, all of which inhibit plant growth and development [23].

Phytohormones in Plant

Recent studies have revealed that different signaling networks, especially those involving phytohormones, ROS, and RNS, arrange many abiotic stress responses [15,24]. Phytohormones are endogenous chemical components that act at the location of synthesis or translocate to different tissues to induce plant growth and enhance crop yield both quantity and quality under different abiotic stress. There are main phytohormones produced in plant tissue (Figure 1).




Figure 1: Main phytohormones produced in plant tissue.

Furthermore, phytohormones are paramount for plants as they use them to regulate their own processes and are implicated in plant interactions with their environment. Phytohormones modulate physiological responses in plant tissues to stimulate adaptation to various abiotic stresses [13].

In recent years, there has been more attention given to the role of phytohormones in parallel with rising salinity stress in different regions worldwide. At salinity stress, various signaling pathways such as phospholipids, plant hormones, and calcium ions are coordinately activated to organize an osmotic modulation or homeostasis regulating plant growth and development [3]. In addition, plant hormonal regulation and calcium ion-dependent modification of enzymatic activities are independently integrated into the stress signaling pathways [5].

Phytohormones play a critical role in plants' capacity to respond to abiotic stresses by regulating a variety of adaptive restraints. They frequently modify gene expression quickly by stimulating or inhibiting the degradation of transcriptional regulators via the ubiquitin-proteasome system [25].

How Plant Hormones Mediate Salt Stress Responses

Rising salinity is considered one of the main abiotic stresses that restrict the growth of numerous crops. Plants developing various adaptation tactics to cope with salinity stress, which including reduce water loss by regulate stomata closing and minimizing growth to maintain plant survival under stress [15,26].

Phytohormones are endogenous substances that regulate plant response and balancing plant growth under salt stress. Phytohormones have an important role in adapting growth and defense trade-offs under stress [27]. There is a significant role for phytohormones in the regulation of plant responses to salinity stress and adaptive plant growth under stress, therefore, exogenous application of phytohormones to plants ameliorates the adverse effect of stress condition [6].

Jasmonates (JA)

Jasmonic acid (JA) is one of the endogenous plant hormones that are synthesized by plants and play a significant role in adaptable plant responses under stress conditions. JA or the related substances (methyl jasmonate, MeJA) are involved in various physiological processes in plants [28].

Under salinity stress, plants synthesize more JA and activate JA signals. Therefore, the exogenous application of JA alleviates abiotic stress tolerance in the plant, it regulates plant growth under salt stress by physiological mechanisms, improving antioxidant enzyme activity, increasing organic solutes such as superoxide dismutase (SOD), peroxidase (POD), catalase (CAT), APX, and chlorophyll B contents, and reducing H₂O₂ concentration, consequently improving plant growth under salinity conditions [29]. JA is involved in varied physiological processes, growth, and development, which include root system formation, fruit production, fruit ripening, enhancing oxidative resistance, and interaction with other phytohormones to increase plant tolerance for abiotic stress [30]. In addition, JA has an important role in plant defense against infection by pests, particularly herbivorous insects and diseases like necrotrophic pathogens [31].

Role of Jasmonic Acid in Plant Under Salinity Stress:

The essential role of JA as a phytohormone to increase plant tolerance and acclimatize abiotic stress is well documented. JA plays numerous roles in plant cells to regulate adaptation to environmental stresses, therefore, under salinity stress, JA levels increase in plant cells and there is more JA signaling [31].

Previous research has shown that exogenous treatments of JA alleviate salinity stress by improving plant antioxidant responses, enhancing seed germination, stimulating plant growth, and increasing the survival rate of seedlings under

abiotic stress Sheteiwy MS, et al. [32], Ghassemi-Golezani, et al. [33] on soybean; Hussain, et al. [34] on rice; Sheyhakinia, et al. [35] on roselle; Ahmad, et al. [36] on maize seedlings.

Furthermore, foliar application of Proline and jasmonic acid enhanced crop yield and improved the fruit quality of the Manzanillo olive tree [37]. In addition, the application of humic acid and jasmonic acid alleviates the adverse effects of salinity on Forage Sorghum by improving antioxidant enzymes [38].

The main effect of JA on plant growth under environmental stresses include:

- Reduce Na cations that enter the cells.
- Enhancing the glycine betaine and soluble protein content.
- Increase antioxidant enzyme activity.
- Improve membrane stability index.
- Increase leaf water content.
- Increase antioxidant enzyme activation.
- Enhancing seed germination.
- Alleviate the negative effects of salinity stress on plant growth.
- Increasing plant acclimation under improper conditions.
- Decreased the amount of herbicide residue.

Cytokinin (CK)

Cytokinin (CK) is one of the main five phytohormones produced in plant tissue, CK has numerous functions in plant metabolism, and it has a significant role in enhancing plant response to various adverse conditions and increasing plant tolerance of both abiotic and biotic stress as well. CKs have the opposing influence of ABA and act as antagonists of ABA in numerous physiological aspects in plants [39]. Cytokinin considered a master phytohormones regulate different processing in plants, particularly cell division, flower bud initiation and differentiation, and delaying senescence. Moreover, CKs synthesis in the root system and transported to the vegetative part through the xylem, under abiotic stress some aerial organs synthesize CK, particularly young tissues, and the influence on root CKs occurs via transport basipetal phloem CK [40]. Cytokinins are vital components in plants for numerous biological functions, like flowering, cell division, seed germination, regulating apical dominance, translocating of nutrients, chloroplast development, senescence, and acclimatization to environmental stresses [41].

Under salinity conditions, the synthesis of cytokinin stimulates metabolisms in plants particularly hormonal and ionic status. Maintaining cytokinin homeostasis is essential for plants experiencing salt stress to enhance plant growth, as well as plant adaptation to abiotic stresses [42].

Furthermore, exogenous application of CKs enhances plant growth and reduces the adverse effect of salinity on the growth and productivity of various crop [33] and Latef, et al. [43] on faba bean; Azadi, et al. [44] on black cumin plants; Feng, et al. [45] on apple seedlings; Sa, et al. [46] on papaya plants). In addition, cytokinin improves the yield crop of rice under abiotic stress conditions by regulating floral primordial activity and modulating rice grain yield [47]. Furthermore, CK stimulates photosynthesis and enhances flower production [48].

Cytokinin-mediated stimulation of ethylene production appears to be of particular importance during water stress. CKs are indispensable for the temporal and differential biosynthesis of SA and JA, which in turn serve as endogenous inducers for distinct classes of PR proteins. CK plays an important role in the regulation of stress-related genes in different plants under adverse environmental conditions.

There is more evidence from previous work that CK has more responses to various stresses. There are important functions for CK in plant responses to biotic stress and improving photosynthesis under salinity stress [49].

In addition, under abiotic stress, Cytokines stimulate ethylene production and have a significant role in the synthesis of endogenous hormones like SA and JA Brenner WG, et al. [50] on Arabidopsis, and Joshi, et al. [47] on Rice. Maintaining biosynthesis and cytokinin balance is very important for plant growth and acclimatization of plants to abiotic stresses. Therefore, CK levels in plants are considered an indicator of salt tolerance particularly in various crops [51].

There is crosstalk with other phytohormones particularly JA, and there is a correlation between Cytokinin and Jasmonic acid. JA inhibits the activity of oxidative enzyme-like cytokinin oxidase (CKX) and cytokinin oxidase/dehydrogenase, consequently, increasing the level of Cytokinin [52].

There are emerging roles of cytokinin as follow:

- Priming agent for meristems growth.
- Regulator of plant Defense growth.
- Required factor against biotic stress.
- Improve photosynthesis.
- Increase flowering of tomato plants.
- Increase seed yield of sensitive rice varieties.

Auxin

It is certainly true that auxin plays numerous roles in plant growth and development. Local concentration gradients of auxin (indole-3-acetic acid [IAA]) are thought to instruct the positioning of organ primordia and stem cell functions and to direct cell division, expansion, and differentiation. It also

plays a role in plant responses to biotic and abiotic stresses. Furthermore, it has been shown that auxin affects several plant stages, including root hair, stem cell formation, and elongation, and has a great and important function in plant growth under salinity stress [53].

It was noticed that the lower levels of active auxin that exist in Arabidopsis mutants exhibit a reduction in plant growth and show enhanced resistance to environmental stresses. Auxin has also been implicated in seed germination under salinity conditions [54].

Numerous references indicate the involvement of auxin in response to salinity stress in plants, but little information is available regarding the mechanisms of salt stress regulation. Moreover, the pathway of auxin signaling is closely related to the early seedling developmental processes of plants [55].

Higher auxin concentrations caused by overexpression of the auxin biosynthetically correlated YUCCA3 (YUC3) gene, which encodes an auxin biosynthetic enzyme, caused hypersensitivity to salinity stress [56]. Interestingly, high salt stress greatly remodeled root architecture by altering auxin accumulation and its redistribution. So, the redistribution of auxin maxima formation in plant tissues is associated with decreased growth [57].

Foliar application of indole-3-acetic acid improved the growth characteristics of salinized fava bean due to increasing the uptake of some nutrients like K^+ , Ca^{2+} , and Mg^{2+} ions, accumulation of free amino acids, soluble sugars, and soluble proteins, as well as the activity of many enzymes such as superoxide dismutase, catalase, peroxidase, and ascorbate peroxidase [58].

Exogenous applications of auxin compounds such as indole-3-acetic acid or indole-3-butyric acid before grafting could promote the growth characteristics of grafted cucumber seedlings and their performance after transplanting in saline conditions [59].

Under high salinity conditions, exogenous application of auxin further suppressed the reduced germination rate of Arabidopsis seeds as compared with control. In contrast, the auxin effects disappeared in the *ntm2-1* mutant. These observations indicate that *ntm2-1* is a molecular link that incorporates auxin signal into salt stress signaling during seed germination, providing a role for auxin in modulating seed germination under high salinity [60].

Foliar application of indole-3-acetic acid (IAA) or 6-benzyl adenine (BA) combined with salt stress, creates a major change in the plant protein profile with various intensities. There are better adaptations and increased saline tolerance,

which was confirmed by the exhibited new protein profiles under both saline and treated with IAA or BA conditions. The IAA-applied plants produced more new protein bands than BA, indicating that IAA has more ability to rescue the faba bean from salt stress and adapt to the salt stress condition [58].

Gibberellins

It is known that gibberellins are plant growth regulators that induce various physiological restraints in plants. Generally, gibberellins are phytohormones implicated in the growth and developmental processes of plants like seed germination, leaf expansion, photo-morphogenesis, stem elongation, and flowering [61].

Plant hormones have a role in regulating the expression of genes, especially abscisic acid, gibberellic acid, and mRNA is strongly induced by high salinity stress and the possible relation between the contents of endogenous GA3 and the acquisition of stress protection [62].

It was noticed that bioactive gibberellins are rapidly decreased when plants are exposed to both biotic and abiotic stress. Moreover, under environmental stress, there is a crosstalk between gibberellin action and other hormones signaling the dominance of plant growth and development. However, the underlying molecular mechanism remains to be demonstrated [63].

Seed germination is one of the plant developmental processes affected by salinity stress. The presence of NaCl during embryogenesis affects the growth balance, moreover, soil salinity suppresses seed germination either by magnificent osmotic stress or by decreasing gibberellin biosynthesis [64].

Under salinity stress, gibberellins act as promoters for plant growth, which can decrease seed dormancy, improve plant gene expression, enhance the synthesis of hydrolase, repair injured cell membranes, and increase seed vitality [65]. Gibberellic acid partially alleviates the felled effect of salinity by increasing wheat vigor, anti-oxidative enzyme activity, and the accumulation of osmolytes [66]. On the contrary,

Application of gibberellin promotes cell division and cell elongation of stressed seedlings and reduces inhibition effects of NaCl, which increases seedling growth on Okra seedlings, moreover, exogenous gibberellic acid application enhanced seedling emergence percentage, seedling growth, and relative water content [67] on Okra seedlings; Yousif, et al. [68] on sorghum. Gibberellic acid significantly affected emergence percentage and alleviated the adverse impact of salinity stress by improving water uptake and increasing cellular membrane plasticity [69], which can stimulate the

activity of amylase in cotyledons and the conversion of insoluble starch into soluble sugars for seed germination and promote radical growth [70].

Furthermore, the application of gibberellin caused an increase in the germination percentage of *Tithonia rotundifolia* Black and pulled off the preventive effects of salinity stress on germination. This may be due to the involvement of GA3 in the activation of cytological enzymes, along with an increase in cell wall plasticity and improved water absorption in papaya plants [69].

The exogenous application with GA3 enhanced seed germination rate under salinity stress. In addition, increasing salt stress levels reduced the growth and grain yield of the crop but increased comparatively with seed treatment with gibberellin [71].

Moreover, gibberellic acid treatments increased the development of plant attributes. The fact that they increase the amino acid content in the embryo of olive cultivars and stimulate the synthesis of hydrolytic enzymes required for the digestion of endospermic starch when seeds renew growth at seed germination [72].

In addition, under salinity stress, the decrease in mineral nutrient uptake may due to Na⁺ induced blockage or reduced activity of the transporters, resulting in an ionic imbalance of K⁺, Ca²⁺, and Mg²⁺ as compared to Na⁺. Whereas, treating two barley cultivars with gibberellin caused a reduction in sodium ion contents [73].

It has been reported that DELLA family proteins, pioneer gibberellin negative regulators, are basically implicated in the modification of environmental signals and other plant hormone signaling pathways. Under abiotic stress, the prominent role of GA in arrangement plant growth is due to DELLA protein-mediated growth restraints on exposure to abiotic stress [74].

Exposure to salinity stress induced a decrease in endogenous levels of bioactive GAs, which coincided with a higher accumulation of DELLA proteins. Moreover, among the different types of plant hormones implicated in the alleviation of deleterious effects on salt stress, gibberellin has been especially important in crops [75]. Moreover, under salt stress, the morphological characteristics of barley were suppressed in all plant growth parameters, while this repression was significantly mitigated after gibberellin application [73].

Proline

Proline is an amino acid with numerous functions, and it is also a signaling molecule that works as a plant growth

regulator by triggering cascade signaling processes. Moreover, proline is the major compound that protects cells by stabilizing proteins and cellular membranes. Proline surpasses as a popular osmolyte in plants and gets up-regulated against various stresses [76]. Thus, many plants synthesize amino acids and amides such as proline, alanine, glutamine, asparagine, and quaternary ammonium bases, as well as different sugars, polyols, and cyclites as a non-specific reactions to stress from excess salinity, drought, or frost [77]. Plants suffering from salinity or drought stress in their root zone respond physiologically by regulating their metabolism to adjust to negative conditions. Therefore, plants accumulate organic solutes particularly low molecular weight materials like proline, betaine, polyols, polyamines, and sugar.

Among the metabolic responses to salt stress, the synthesis of compatible osmolytes and non-toxic organic compounds is responsible for the osmotic equilibrium between the cytoplasm and various cell compartments. In many plants, free proline accumulates in response to the prescription of a wide range of biotic and abiotic stresses. Proline controls the expression of a number of genes related to antioxidant enzymes in salt conditions. One gene, 1-pyrroline-5-carboxylate synthase, is responsible for up-regulating the stress-stimulated accumulated of proline under salt stress [78].

Most efforts have been made to consideration of the ability of proline to mediate osmotic adjustment, stabilize subcellular structures, and scavenge toxic oxygen derivatives. High levels of proline synthesized during stress conditions also preserve the NAD (P)⁺/NAD(P)H ratio [3].

Proline biosynthesis, degeneration, and its accumulation in plants are regulated by various abiotic stresses and salinity has great importance [79].

There is a negative relationship between salt tolerance and proline accumulation. Consequently, accumulation of proline in *Pancreaticum maritimum* L. may contribute to cellular adaptation to salt stress or be the result of metabolic changes caused by salinity [80]. Furthermore, it was noticed that proline biosynthesis happened because of the disturbance in cell homeostasis or fluctuation in the use of photosynthesis leftover for proline biosynthesis at the expense of plant growth [81].

Plants have developed various saving mechanisms to maintain normal cellular metabolism and prohibit cellular constituent damage including the accumulation of ions and osmolytes such as proline [82].

One of the efficient saving mechanisms of plants against hyperosmotic stress is the increased endogenous level of

compatible solutes like proline, ectoine, glycine betaine, and sorbitol, in addition, there is a positive correlation between the endogenous levels of proline and increasing salinity levels [42].

Proline provides tolerance against different abiotic stresses by increasing their endogenous levels and their intermediate enzymes in plants, therefore, the exogenous application of proline increases the endogenous level of proline in bean (*Phaseolus vulgaris* L.) [83].

Many plants accumulate a higher level of proline in contrast to other amino acids when exposed to the high salt content in the soil. When the salt has been accumulated in the vacuole proline acts as a mediator of osmotic adjustment stabilizing, to save cell membranes, several enzymes, and metabolic machinery [84].

Moreover, when plants are subjected to saline conditions, proline plays a very important role in preserving their growth. Therefore, proline contributes to osmotic adjustment, stabilization, and saving the membrane integrity and macromolecules from the harmful effects of salinity and as a hydroxyl radical scavenger. Furthermore, it was suggested that, in saline conditions, proline accumulation in leaves occurred to protect chlorophyll content and turgor to save the photosynthetic activity [85].

Moreover, exogenous application of proline enhances the crop tolerance against different abiotic stresses especially salinity by protecting them from the severe effects of ROS [42]. Foliar application of proline at the rate of 100 mg.l increased plant height, the number of branches as well as fresh and dry weights of leaves of *Beta vulgaris* L. [86]. Exogenous application of proline at the rate of 25 mM increases in *Vicia faba* growth characters [87]. All proline levels (3, 6, or 9 mM) caused a significant increase in shoot length, number of leaves and branches per plant, and shoot fresh and dry weights per plant of both lupine varieties compared with control [88].

In addition, plants resort to promoting their endogenous level of proline with continuously increasing levels of salinity [89]. Moreover, the exogenous application of proline mitigates the opposite effects of salt by decreasing the accumulation of Na⁺ and Cl⁻ in plants [90].

Conclusion

Plant growth regulators such as auxin and gibberellin improve salt tolerance and mitigate salt stress by stimulating plant growth and development. Recent advances in the research, which is dependent on protein profile, indicated that the disappearance of polypeptide bands in response to

stress and foliar applications of phytohormones such as IAA showed a new protein band and newly expressed protein that might diminish the adverse effects of salt stress and increase better adaptation to plants. Moreover, under salinity stress, exogenous application of gibberellins promotes plant growth by improving water uptake, decreasing seed dormancy, activating antioxidant enzymes, and improving plant gene expression. All of these alleviated the negative effects of salinity stress on plants.

In addition, salt stress imposes severe effects on plant growth and productivity by interrupting normal metabolic processes and proline may mitigate the adverse impact of salt by reducing osmotic stress that consequently safeguards the membrane integrity and its function. The exogenous application of proline could offer a simple and economical approach for farmers to decrease the crop loss risk in salt contaminated land. Otherwise, more studies are needed at physiological and molecular levels to gain deeper prudence in understanding the interaction of NaCl persuades oxidative stress and mitigation mechanism of exogenous proline in plants.

References

1. Nykiel M, Gietler M, Fidler J, Prabucka B, Graska J, et al. (2022) Signal transduction in cereal plants struggling with environmental stresses: from perception to response. *Plants* 11(8): 1009[]]
2. Mustafa G, Akhtar MS, Abdullah R (2019) Global Concern for Salinity on Various Agro-Ecosystems. In: Akhtar M (Ed.), *Salt Stress, Microbes, and Plant Interactions: Causes and Solution*. Springer, Singapore, 1: 1-19[]]
3. Abobatta WF (2020) Plant Responses and Tolerance to Combined Salt and Drought Stress. In: Tanveer M, (Ed.), *Salt and Drought Stress Tolerance in Plants. Signaling and Communication in Plants*. Springer, Cham, pp: 17-52.
4. Riaz M, Arif MS, Ashraf MA, Mahmood R, Yasmeen T, et al. (2019) A comprehensive review on rice responses and tolerance to salt stress. In: Hasanuzzaman M (Ed.), *Advances in rice research for abiotic stress tolerance*. Woodhead Publishing, United Kingdom, pp: 133-158[]]
5. Ryu H, Cho YG (2015) Plant hormones in salt stress tolerance. *Journal of Plant Biology* 58(3): 147-155[]]
6. Sadiq Y, Zaid A, Khan MMA (2020) Adaptive Physiological Responses of Plants under Abiotic Stresses: Role of Phytohormones. In: Hasanuzzaman M (Ed.), *Plant Ecophysiology and Adaptation under Climate Change: Mechanisms and Perspectives I*. Springer, Singapore, pp: 797-824[]]
7. Pratima P, Chawla W (2019) Influence of plant growth regulators on growth and yield of pome and stone fruits. *Journal of Pharmacognosy and Phytochemistry* 8(1S): 557-565[]]
8. Diego ND, Spíchal L (2020) Use of plant metabolites to mitigate stress effects in crops. In: Xu L (Ed.), *The chemical biology of plant biostimulants*. Wiley, New Jersey, United States, pp: 261-300[]]
9. Khan N, Bano A, Ali S, Babar MA (2020) Crosstalk amongst phytohormones from planta and PGPR under biotic and abiotic stresses. *Plant Growth Regulation* 90(2): 189-203.
10. Jiroutova P, Oklestkova J, Strnad M (2018) Crosstalk between brassinosteroids and ethylene during plant growth and under abiotic stress conditions. *International journal of molecular sciences* 19(10): 3283.
11. Bhatt D, Nath M, Sharma M, Bhatt MD, Bisht DS, et al. (2020) Role of growth regulators and phytohormones in overcoming environmental stress. In: Tripathi DK (Eds.) *Protective Chemical Agents in the Amelioration of Plant Abiotic Stress: Biochemical and Molecular Perspectives*. Wiley, New Jersey, United States, pp: 254-279[]]
12. Berwal MK, Kumar R, Prakash K, Rai GK, Hebbar KB (2021) Antioxidant defense system in plants against abiotic stress. In: Bagati S, et al. (Eds.) *Abiotic stress tolerance mechanisms in plants*. CRC Press, Taylor & Francis, London, UK, pp: 175-202[]]
13. EL Sabagh A, Islam MS, Hossain A, Iqbal MA, Mubeen M, et al. (2022) Phytohormones as growth regulators during abiotic stress tolerance in plants. *Frontiers in Agronomy* 4: 765068[]]
14. Hannachi S, Steppe K, Eloudi M, Mechi L, Bahrini I, et al. (2022) Salt stress induced changes in photosynthesis and metabolic profiles of one tolerant ('Bonica') and one sensitive ('Black beauty') eggplant cultivars (*Solanum melongena* L.). *Plants* 11(5): 590[]]
15. Abobatta WF (2020) Plant Responses and Tolerance to Extreme Salinity: Learning from Halophyte Tolerance to Extreme Salinity. In: Tanveer M (Ed.), *Salt and Drought Stress Tolerance in Plants*. Springer, USA, pp: 177-210.
16. Ziogas V, Tanou G, Morianou G, Kourgialas N (2021) Drought and salinity in citriculture: optimal practices to alleviate salinity and water stress. *Agronomy* 11(7): 1283[]]
17. Malakar P, Chattopadhyay D (2021) Adaptation of plants to salt stress: the role of the ion transporters. *Journal of*

- Plant Biochemistry and Biotechnology 30: 668-683[]]
18. Bidalia A, Vikram K, Yamal G, Rao KS (2019) Effect of salinity on soil nutrients and plant health. *Salt Stress, Microbes, and Plant Interactions: Causes and Solution* 1: 273-297[]]
 19. Atta K, Mondal S, Gorai S, Singh AP, Kumari A, et al. (2023). Impacts of salinity stress on crop plants: improving salt tolerance through genetic and molecular dissection. *Frontiers in Plant Science* 14: 1241736[]]
 20. Zhao C, Zhang H, Song C, Zhu JK, Shabala S (2020) Mechanisms of plant responses and adaptation to soil salinity. *The innovation* 1(1): 1-41.
 21. Hasanuzzaman M, Raihan MRH, Masud AAC, Rahman K, Nowroz F, et al. (2021) Regulation of reactive oxygen species and antioxidant defense in plants under salinity. *International Journal of Molecular Sciences* 22(17): 9326[]]
 22. Pan T, Liu M, Kreslavski VD, Zharmukhamedov SK, Nie C, et al. (2021) Non-stomatal limitation of photosynthesis by soil salinity. *Critical Reviews in Environmental Science and Technology* 51(8): 791-825[]]
 23. Hnilickova H, Kraus K, Vachova P, Hnilicka F (2021) Salinity stress affects photosynthesis, malondialdehyde formation, and proline content in *Portulaca oleracea* L. *Plants* 10(5): 845[]]
 24. Sharma P, Sharma P, Arora P, Verma V, Khanna K, et al. (2019) Role and regulation of ROS and antioxidants as signaling molecules in response to abiotic stresses. In: Khan NA (Ed.), *Plant signaling molecules*. Woodhead Publishing, Sawston, UK, pp: 141-156[]]
 25. Saharan BS, Brar B, Duhan JS, Kumar R, Marwaha S, et al. (2022) Molecular and physiological mechanisms to mitigate abiotic stress conditions in plants. *Life* 12(10): 1634[]]
 26. Cao H, Ding R, Kang S, Du T, Tong L, et al. (2023) Drought, salt, and combined stresses in plants: effects, tolerance mechanisms, and strategies. *Advances in Agronomy* 178: 107-163[]]
 27. Yu Z, Duan X, Luo L, Dai S, Ding Z, et al. (2020) How plant hormones mediate salt stress responses. *Trends in plant science* 25(11): 1117-1130.
 28. Rehman M, Saeed MS, Fan X, Salam A, Munir R, et al. (2023) The Multifaceted Role of Jasmonic Acid in Plant Stress Mitigation: An Overview. *Plants* 12(23): 3982[]]
 29. Siddiqi KS, Husen A (2019) Plant response to jasmonates: current developments and their role in changing environment. *Bulletin of the National Research Centre* 43(1): 1-11.
 30. Ahmad B, Raina A, Naikoo MI, Khan S (2019) Role of methyl jasmonates in salt stress tolerance in crop plants. In: Khan MIR, (Eds.) *Plant Signaling Molecules*, Wood head Publishing, Sawston, United Kingdom, pp: 371-384[]]
 31. Wang Y, Mostafa S, Zeng W, Jin B (2021) Function and mechanism of jasmonic acid in plant responses to abiotic and biotic stresses. *International Journal of Molecular Sciences* 22(16): 8568[]]
 32. Sheteiwy MS, Shao H, Qi W, Daly P, Sharma A, et al. (2021) Seed priming and foliar application with jasmonic acid enhance salinity stress tolerance of soybean (*Glycine max* L.) seedlings. *Journal of the Science of Food and Agriculture* 101(5): 2027-2041[]]
 33. Ghassemi-Golezani K, Samea-Andabjadid S (2022) Exogenous cytokinin and salicylic acid improve amino acid content and composition of faba bean seeds under salt stress. *Gesunde Pflanzen* 74(4): 935-945[]]
 34. Hussain S, Zhang R, Liu S, Li R, Wang Y, et al. (2022) Methyl Jasmonate Alleviates the Deleterious Effects of Salinity Stress by Augmenting Antioxidant Enzyme Activity and Ion Homeostasis in Rice (*Oryza sativa* L.). *Agronomy* 12(10): 2343[]]
 35. Sheyhakinia S, Bamary Z, Einali A, Valizadeh J (2020) The induction of salt stress tolerance by jasmonic acid treatment in roselle (*Hibiscus sabdariffa* L.) seedlings through enhancing antioxidant enzymes activity and metabolic changes. *Biologia* 75: 681-692[]]
 36. Chen YE, Mao JJ, Sun LQ, Huang B, Ding CB, et al. (2021) Exogenous application of melatonin induces tolerance to salt stress by improving the photosynthetic efficiency and antioxidant defense system of maize seedling. *Journal of Plant Growth Regulation* 40(3): 1270-1283[]]
 37. El-Sayed OM, El-Gammal OHM, Salama ASM (2014) Effect of ascorbic acid, proline and jasmonic acid foliar spraying on fruit set and yield of Manzanillo olive trees under salt stress. *Scientia Horticulturae* 176: 32-37.
 38. Ali AYA, Ibrahim MEH, Zhou G, Nimir NEA, Jiao X, et al. (2019) Ameliorative effects of jasmonic acid and humic acid on antioxidant enzymes and salt tolerance of forage sorghum under salinity conditions. *Agronomy Journal* 111(6): 3099-3108.
 39. Joshi V, Sahu A, Joshi N, Vyas A, Shah K, et al. (2023) Signaling crosstalk between cytokinins and abscisic

- acid in plant defense, growth, and development. In: Zhang W (Ed.), Hormonal Cross-Talk, Plant Defense and Development. Cambridge, Massachusetts, United States, pp: 149-170.]
40. Kaur R, Sharma N, Tikoria R, Ali M, Kour S, et al. (2022) Insights into Biosynthesis and Signaling of Cytokinins During Plant Growth, Development and Stress Tolerance. In: Aftab T, (Eds.) Auxins, Cytokinins and Gibberellins Signaling in Plants. Signaling and Communication in Plants. Springer, Germany, pp: 153-187.
 41. Wang Y, Shen W, Chan Z, Wu Y (2015) Endogenous Cytokinin Overproduction Modulates ROS Homeostasis and Decreases Salt Stress Resistance in Arabidopsis Thaliana. Front Plant Sci 6:1004.
 42. Nounjan N, Nghia PT, Theerakulpisut P (2012) Exogenous proline and trehalose promote recovery of rice seedlings from salt-stress and differentially modulate antioxidant enzymes and expression of related genes. J Plant Physiol 169(6): 596-604.
 43. Latef AAA, Hasanuzzaman M, Tahjib-Ul-Arif M (2021) Mitigation of salinity stress by exogenous application of cytokinin in faba bean (*Vicia faba* L.). Notulae Botanicae Horti Agrobotanici Cluj-Napoca 49(1): 12192-12192.]
 44. Azadi F, Hatami A, Salek Mearaji H (2022) The effect of cytokinin foliar on morpho-physiological traits, yield and yield components of black cumin (*Nigella sativa* L.) under salinity stress conditions. Environmental Stresses in Crop Sciences 15(4): 975-990.]
 45. Feng Y, Liu J, Zhai L, Gan Z, Zhang G, et al. (2019) Natural variation in cytokinin maintenance improves salt tolerance in apple rootstocks. Plant Cell Environ 42(2): 424-436.
 46. Sá FVDS, Brito ME, Silva LDA, Moreira RC, Paiva EPD, et al. (2020) Exogenous application of phytohormones mitigates the effect of salt stress on Carica papaya plants. Revista Brasileira de Engenharia Agrícola e Ambiental 24: 170-175.]
 47. Joshi R, Sahoo KK, Singh AK, Anwar K, Pundir P, et al. (2020) Enhancing trehalose biosynthesis improves yield potential in marker-free transgenic rice under drought, saline, and sodic conditions. Journal of Experimental Botany 71(2): 653-668.]
 48. Janowska B, Andrzejak R (2022) Cytokinins and gibberellins stimulate the flowering and post-harvest longevity of flowers and leaves of Calla lilies (*Zantedeschia Spreng.*) with colourful inflorescence spathes. Agronomy 12(8): 1859.]
 49. Yu Y, Li Y, Yan Z, Duan X (2021) The role of cytokinins in plant under salt stress. Journal of Plant Growth Regulation 41(6): 1-13.]
 50. Brenner WG, Schmölling T (2015) Summarizing and exploring data of a decade of cytokinin-related transcriptomics. Frontiers in plant science 6: 29.
 51. Ashraf M, Munns R (2022) Evolution of approaches to increase the salt tolerance of crops. Critical Reviews in Plant Sciences 41(2): 128-160.]
 52. Avalbaev A, Allagulova C, Maslennikova D, Fedorova K, Shakirova F (2020) Methyl jasmonate and cytokinin mitigate the salinity-induced oxidative injury in wheat seedlings. Journal of Plant Growth Regulation 40(4): 1741-1752.
 53. Mazzoni-Putman SM, Brumos J, Zhao C, Alonso JM, Stepanova AN (2021) Auxin interactions with other hormones in plant development. Cold Spring Harbor Perspectives in Biology 13(10): a039990.]
 54. Abdel Latef AAH, Tahjib-Ul-Arif M, Rhaman MS (2021) Exogenous auxin-mediated salt stress alleviation in faba bean (*Vicia faba* L.). Agronomy 11(3): 547.]
 55. Gomes GLB, Scortecci KC (2021) Auxin and its role in plant development: structure, signalling, regulation and response mechanisms. Plant Biology 23(6): 894-904.]
 56. Cao X, Yang H, Shang C, Ma S, Liu L, et al. (2019) The roles of auxin biosynthesis YUCCA gene family in plants. International Journal of Molecular Sciences 20(24): 6343.]
 57. Smolko A, Bauer N, Pavlović I, Pěnčík A, Novák O, et al. (2021) Altered root growth, auxin metabolism and distribution in Arabidopsis thaliana exposed to salt and osmotic stress. International Journal of Molecular Sciences 22(15): 7993.]
 58. Abdel Latef AAH, Akter A, Tahjib-Ul-Arif M (2021) Foliar Application of auxin or cytokinin can confer salinity stress tolerance in *Vicia faba* L. Agronomy 11(4): 790.
 59. Balliu A, Sallaku G (2017) Exogenous auxin improves root morphology and restores growth of grafted cucumber seedlings. Hort Sci (Prague) 44(2): 82-90.
 60. Albertos P, Tatematsu K, Mateos I, Nakabayashi K, Nambara E, et al. (2021) Redox feedback regulation of ANAC089 signaling alters seed germination and stress response. Cell Reports 35(11): 109263.]
 61. Chauhan A, Rajput N, Kumar D, Kumar A, Chaudhry AK (2016) Effect of different salt concentration seed

- germination and seedling growth of different varieties of oat (*Avena sativa* L.). *Int J Inform Res Rev* 3(7): 2627-2632.
62. Ahmad I, Zhu G, Zhou G, Song X, Hussain S, et al. (2022) Pivotal role of phytohormones and their responsive genes in plant growth and their signaling and transduction pathway under salt stress in cotton. *International Journal of Molecular Sciences* 23(13): 7339.
 63. Castro-Camba R, Sánchez C, Vidal N, Vielba JM (2022) Interactions of gibberellins with phytohormones and their role in stress responses. *Horticulturae* 8(3): 241.
 64. Soni PG, Basak N, Rai AK, Sundha P, Chandra P, et al. (2023) Occurrence of Salinity and Drought Stresses: Status, Impact, and Management. In: Kumar A (Ed.), *Salinity and Drought Tolerance in Plants*. Springer, Singapore, pp: 1-28.
 65. Kandil AA, Sharief AE, Abido WAE, Awed AM (2014) Effect of gibberellic acid on germination behaviour of sugar beet cultivars under salt stress conditions of Egypt. *Sugar Tech* 16(2): 211-221.
 66. Neelambari, Mandavia C, Ganesh S (2018) Curative effect of ascorbic acid and gibberellic acid on wheat (*Triticum aestivum* L.) metabolism under salinity stress. *Int J Curr Microbial App Sci* 7(1): 522-533.
 67. Wang YH, Zhang G, Chen Y, Gao J, Sun YR, et al. (2019) Exogenous application of gibberellic acid and ascorbic acid improved tolerance of okra seedlings to NaCl stress. *Acta Physiologiae Plantarum* 41(6): 1-10.
 68. Ali AYA, Ibrahim MEH, Zhou G, Nimir NEA, Elsiddig AMI, et al. (2021) Gibberellic acid and nitrogen efficiently protect early seedlings growth stage from salt stress damage in sorghum. *Scientific Reports* 11(1): 6672.
 69. Lay P, Basvaraju GV, Sarika G, Amrutha N (2013) Effect of seed treatments to enhance seed quality of papaya (*Carica papaya* L.) cv. Surya. *Glob J Biol Agric Health Sci* 2(3): 221-225.
 70. Ibrahim MEH, Zhu X, Zhou G, Yousif AAA, Mohammed AIE, et al. (2019) Response of some wheat varieties to gibberellic acid under saline conditions. *Agro systems Geosciences Environ* 2(1): 1-7.
 71. Javid MG, Sorooshzadeh A, Moradi F, Sanavy S, Allahdadi I (2011) The role of phytohormones in alleviating salt stress in crop plants. *Aust J Crop Sci* 5(6): 726-734.
 72. Shekafandeh A, Smoushtaghi, Sirooenejad E (2017) Influence of gibberellin on increasing of sodium chloride tolerance via some morpho-physiological changes in two olive cultivars. *Agri Conspectus Sci* 82(4): 367-373.
 73. Amal MEA, Heba IM (2014) The effect of the exogenous gibberellic acid on two salt stressed barley cultivars. *European Scientific Journal* 10(6): 1857-7881.
 74. Huang X, Tian H, Park J, Oh DH, Hu J, et al. (2023) The master growth regulator DELLA binding to histone H2A is essential for DELLA-mediated global transcription regulation. *Nature plants* 9(8): 1291-1305.
 75. Colebrook EH, Thomas SG, Phillips AL, Hedden P (2014) The role of gibberellin signalling in plant responses to abiotic stress. *J Exp Biol* 217(1): 67-75.
 76. Meena M, Divyanshu K, Kumar S, Swapnil P, Zehra A, et al. (2019) Regulation of L-proline biosynthesis, signal transduction, transport, accumulation and its vital role in plants during variable environmental conditions. *Heliyon* 5(12): e02952.
 77. Ingrisano R, Tosato E, Trost P, Gurrieri L, Sparla F (2023) Proline, Cysteine and Branched-Chain Amino Acids in Abiotic Stress Response of Land Plants and Microalgae. *Plants* 12(19): 3410.
 78. Kim GB, Nam YW (2013) A novel $\Delta(1)$ -pyrroline-5-carboxylate synthetase gene of *Medicago truncatula* plays a predominant role in stress-induced proline accumulation during symbiotic nitrogen fixation. *J plant Physiol* 170(3): 291-302.
 79. Shafi A, Zahoor I, Mushtaq U (2019) Proline accumulation and oxidative stress: Diverse roles and mechanism of tolerance and adaptation under salinity stress. *Salt Stress, Microbes, and Plant Interactions: Mechanisms and Molecular Approaches* 2: 269-300.
 80. Carfagna S, Salbitani G, Innangi M, Menale B, De Castro O, et al. (2021) Simultaneous biochemical and physiological responses of the roots and leaves of *Pancreaticum maritimum* (Amaryllidaceae) to mild salt stress. *Plants* 10(2): 345.
 81. Silambarasan N, Natarajan S (2014) Biochemical responses of Sankankuppi (*Clerodendron inerme* L.) to salinity stress. *African Journal of Agricultural Research* 9(15): 1151-1160.
 82. Kolupaev YE, Yastreb TO, Oboznyi AI, Ryabchun NI, Kirichenko VV (2016) Constitutive and cold-induced resistance of rye and wheat seedlings to oxidative stress. *Russian Journal of Plant Physiology* 63(3): 326-337.
 83. Aggarwal M, Sharma S, Kaur N, Pathania D, Bhandhari K, et al. (2011) Exogenous proline application reduces phytotoxic effects of selenium by minimising oxidative

- stress and improves growth in bean (*Phaseolus vulgaris* L.) seedlings. *Biol Trace Elem Res* 140(3): 354-367.
84. Zadehbagheri M, Azarpanah A, Javanmardi S (2014) Proline metabolite transport an efficient approach in corn yield improvement as response to drought conditions. *International Journal of Farming and Allied Sciences* 3(5): 453-461.
 85. Liu D, Ford KL, Roessner U, Natera S, Cassin AM, et al. (2013) Rice suspension cultured cells are evaluated as a model system to study salt responsive networks in plants using a combined proteomic and metabolomic profiling approach. *Proteomics* 13(12&13): 2046-2062.
 86. El-Sherbeny MR, Da Silva JAT (2013) Foliar treatment with proline and tyrosine affect the growth and yield of beetroot and some pigments in beetroot leaves. *J Hort Res* 21(2): 95-99.
 87. Dawood MG, Taie HAA, Nassar RMA, Abdelhamid MT, Schmidhalter U (2014) The changes induced in the physiological, biochemical and anatomical characteristics of *Vicia faba* by the exogenous application of proline under seawater stress. *South Afr J Bot* 93: 54-63.
 88. Rady MM, Talaat NB, Abdelhamid MT, Shawky BT, Desoky ESM (2019) Maize (*Zea mays* L.) grains extract mitigates the deleterious effects of salt stress on common bean (*Phaseolus vulgaris* L.) growth and physiology. *The Journal of Horticultural Science and Biotechnology* 94(6): 777-789.
 89. Molazem D, Qurbanov EM, Dunyamaliyev SA (2010) Role of proline, Na and chlorophyll content in salt tolerance of corn (*Zea mays* L.). *Amer Eur J Agric Environ Sci* 9(3): 319-324.
 90. Khattab EA, Afifi MH (2009) Effect of proline and glycinebetain on canola plants grown under salinity stress condition. *Mod J Appl Biol Sci Crop Sci* 3(11): 42-51.