

Suboptimal Sleep in Adolescents: Socio-Cultural Risk Factors to Consider

Leigh Miles Jackson*

The National Academies of Sciences, Engineering and Medicine, USA

***Corresponding author:** Dr. Leigh Miles Jackson, Ph.D, The National Academies of Sciences, Engineering, and Medicine. 500 Fifth Street, NW Washington, DC 20001, USA, Tel no: 202-334-2047; Email: LMJackson@nas.edu

Received Date: August 03, 2018; **Published Date:** August 14, 2018

Keywords: Suboptimal; Sleep in adolescents; Risk factors

Abbreviations: CDC: Centers for Disease Control and Prevention; SES: Socioeconomic Status

Introduction

Chronic sleep problems, such as sleep loss and disrupted sleep, are associated with poor health outcomes and continue to contribute to the nation's public health burden. Growing incidence of sleep problems among adolescents suggests that the nation's youth may be at increased risk for short- and long-term health consequences. The intent of this brief commentary is to increase public awareness of the complex context that shapes the quality and quantity of adolescent sleep, to underscore the conclusion that adolescence is a particularly salient developmental period for prevention and intervention research, and to highlight the findings that adolescents of color may be at particularly high risk for negative health outcomes related to suboptimal sleep.

Sleep and Adolescents

Roughly 70 million Americans suffer from chronic sleep problems [1]. Suboptimal sleep not only affects an

individual's physical safety and quality of life, but is also linked to a variety of psychiatric and somatic disorders (e.g. depression, obesity), elevated stress, and increased mortality [2,3]. Although the precise mechanisms underlying the complex relation between mammalian sleep and health outcomes are unclear, the importance of studying sleep patterns and behavior remain widely appreciated [4,5].

Adults with sleep problems commonly report that their sleep disturbances began in adolescence [4,6]. This is not particularly surprising, in that sleep profiles naturally change across each major developmental period [7,8]. Adolescence, however, is a particularly vulnerable period during which youth experience significant changes in sleep patterns and behaviors, reductions in total sleep time, and instances of sleep disruption (e.g., difficulty falling asleep, difficulty staying asleep) [8-10]. These changes in sleep are often attributed to later bedtimes, partially resulting from decreased parental supervision, more societal demands (e.g., early school start times, increased academic requirements), and increased stress (e.g., interpersonal difficulties, academic demands) [11,12].

In adolescents, insufficient amounts of sleep is linked to poorer overall health, altered sleep architecture, greater day time sleepiness, increased risk-taking activities, and decreased emotion regulation, as compared to those

receiving adequate sleep [9,10]. Furthermore, sleep loss and the associated deficits in emotion regulation is linked to the predisposition for, and increased severity of, internalizing and externalizing disorders (e.g. depression, aggression) that may persist well in to adulthood [9,13,14].

An important step toward the intervention and prevention of health consequences related to poor sleep is a greater understanding of the social, cultural, and environmental factors that impact sleep behavior during the particularly malleable period of adolescence. Below is a brief overview of the impact of five external factors on adolescent sleep, as well as a short discussion on the potential for race/ethnicity and socioeconomic environments to moderate the relation between sleep and health outcomes.

External Factors that Impact Adolescent Sleep

Electronic media and technology

The integral role of electronic media and technology in the lives of adolescents has increased over the last few decades [15,16]. And unfortunately, this increased consumption has contributed to a disruption in sleep patterns and behaviors. In fact, recent reviews conclude that prolonged use of media and technology is associated with delayed bedtimes, longer sleep onset latency (taking longer to fall asleep), and shorter overall sleep time in children and adolescents [17,18].

These findings make sense, in that late night use of media and technology opposes the biological processes of circadian timing and sleep pressure, and contributes to states of wakefulness and arousal, largely as are salt of sustained light exposure. When activated, a vast number of media devices emit steady waves of blue light [19,20]. The mammalian circadian system is maximally sensitive to the blue spectrum of light (peaksensitivity~460nm) [21]. Blue light exposure during night hours can suppress the secretion of the “sleep hormone,” melatonin [22,23]. And exacerbate sleep delay. Moreover, night time exposure to light can increase core body temperature and heart rate and enhance acute alertness [24,25].

Interestingly, evidence suggests that adolescents’ relationship with sleep and media and technology can be bi-directional, meaning that adolescents with sleep problems may also seek out these electric devices to serve as a sleep aid [10]. Regardless of the direction of the relationship, studies suggest that with chronic repetition, seemingly unimportant habits or behaviors at bedtime can influence outcomes of sleep and have detrimental consequences on an individual’s health and well-being

[26].

Caffeine

Caffeine is a known stimulant and energy enhancer. Over the last few decades, there has been a steady increase in the intake of caffeinated drinks throughout the day and into the evening among youth [15,27,28]. Unfortunately, the “coolness factor” of energy drinks and coffee shop sippy cups has simply become the new norm. Not surprisingly, caffeine impacts sleep. Caffeine is a central nervous system stimulant that increases then euro transmission of excitatory factors (e.g. glutamate, epinephrine) and results in feelings of wakefulness and alertness [29]. Moderate doses of caffeine advance REM sleep, reduce overall slow-wave sleep, and disrupt sleep continuity [27,30]. Moreover, there is a 2-fold increase in sleep disturbances and a significant decrease in over all sleep time in adolescents who reported high caffeine intake, as compared to those whose intake levels were low [15,28].

Cigarette smoking

The Surgeon General reports that nearly 90 percent of adult smokers started smoking by age 18 [31]. In addition, since 2011, there has been a marked increase in the use of e-cigarette among middle and high school students [32]. These statistics on use suggest that adolescents are a vulnerable population for the short- and long-term effects of nicotine on sleep patterns and behavior. A side from the well-known associations with cancer, cigarette smoking is also tightly linked to deficits in sleep quality and duration in adults [33]. The nicotine within cigarettes activates a wide range of cholinergic-driven processes, the most relevant to this discussion is the cholinergic regulation of sleep architecture (e.g., REM stage) [34,35]. Individuals who smoke display lower levels of sleep quality (e.g. more sleep apneas, leg movements, less restorative sleep) and significant changes in sleep architecture (shorter sleep period time, longer sleep latency, higher rapid eye movement (REM) sleep density), as compared to healthy non-smokers [34,36]. The chronic ingestion of nicotine may be, in part, one mechanism linking smoking to the observed changes in sleep patterns.

It remain sun clear as to whether direct exposure to nicotine (or another particular product within the tobacco) disrupts optimal sleep, but studies using only nicotine patches on healthy nonsmokers reveal disruptions in sleep behavior (e.g. decreased total sleep time, reduction in REM sleep) similar to those in sleep studies with smokers [37,38]. Thus, nicotine usage, or possibly the overnight withdraw all from nicotine, may contribute to suboptimal sleep [39,40]. These are notable

findings when considering the short- and long-term effects of nicotine in youth populations.

Alcohol

Adolescence is a unique developmental period characterized by growing behavioral autonomy, decreasing parental supervision, and increasing access to, and exploration of, risk-taking activities, such as alcohol consumption [41]. According to reports from the Centers for Disease Control and Prevention (CDC), alcohol is the most commonly abused drug by youths within the United States, above recent rates of tobacco usage [42]. Alcohol, a depressive drug that activates the inhibitory GABA system, impairs alertness during the day and alters sleep profiles at night [43,44]. Empirical tests on the effects of alcohol use and abuse on adolescent sleep are limited; therefore, researchers largely rely on subjective surveys and adult studies. Resoundingly, sleep studies in adults report that occasional alcohol use may improve sleep [45]. However, moderate excessive use of alcohol increases the latency to fall asleep, produces a dose-dependent alteration in REM sleep, and disrupts sleep continuity (increased waking periods), particularly during the second half of nightly sleep, following the metabolism of the ingested alcohol [45,46]. Given the extent of alcohol use among youth, the relation between alcohol use and sleep problems is an important public health issue with wide spread health implications.

Obesity

For youth (ages 2-19), rates of obesity have increased over the last 20 years. Recent data from the CDC's National Health and Nutrition Examination Survey estimates that roughly 20 percent of youth are obese. Also in these data, are striking disparities in obesity rates across race/ethnicity? For example, 14 percent of non-Hispanic White youths are obese, as compared to roughly 22 percent of African-American youths and roughly 26 percent of Hispanic youth [47]. Given these recent trends (particularly for youths of color), there have been increased concerns regarding the associations between weight, sleep and health.

Studies in adolescents, using both subjective and objective measures, demonstrate a negative association between obesity and sleep duration and deficiency [48,49]. The association between sleep loss and the high risk for obesity is thought to be due to the effects of sleep debt on normative metabolic processes [50]. The secretion of metabolic factors (e.g., glucose, leptin, ghrelin) - important in appetite and feeding behavior-follow distinct circadian rhythm profiles that are sensitive to levels of sleep loss or sleep deprivation [51].

For example, sleep loss alters night time insulin and glucose profiles, promotes increases in ghrelin (an appetite-stimulating hormone), and reduces glucose tolerance and leptin (a satiety-stimulating hormone) levels [52,53]. Additionally, following sleep restriction in healthy young adults, Spiegel et al. [51] found decreases in glucose clearance, lower insulin responses to glucose, decreased carbohydrate tolerance, and higher glucose levels after eating, as compared to individuals for whom sleep time was extended. Furthermore, these metabolic changes contribute to an increased preference for high caloric foods, increased food intake, changes in day time energy use, and are well-recognized precursors of obesity or insulin resistance (type-2diabetes) [54,55].

On the other hand, obesity also identified as an important risk factor for sleep loss [56]. Sleep-related breathing disorders (e.g., sleep apnea) are common in over weight individuals, and may be due to a compromised respiratory structure and/or function [57]. Patients with sleep apnea often report difficulty falling asleep, staying asleep, waking un refreshed, and decreased day time physical activity. Importantly although inadequate sleep is not the sole cause of obesity [58]. Evidence indicates that weight loss can reduce episodes of sleep-related apnea and improve health outcomes [59]. Thus, identifying risk factors that contribute to suboptimal sleep can help guide the construction of early intervention programs that aim to ameliorate or prevent the onset of associated health problems.

Potential Modulators of the Relation Between Adolescent sleep and health outcomes

Adolescents of a particular race, ethnic background, or socioeconomic status (SES) may be at especially high risk for the development of suboptimal sleep and negative health outcomes, as compared to their peers. Here, we highlight the importance of considering race/ethnicity and SES when examining the relation between sleep and health outcomes. Race is a complex term that describes a categorization of people. There is ongoing debate concerning the relation of biological race to health outcomes [e.g.,74], but in this commentary, the complex term "race" refers to a self-identified cultural group that is not based on a particular genetic makeup. The construct of SES is a marker of location within the social structure or class and is composed of a number of variables including property ownership, occupation, education, income, number of house hold occupants, and marital status. It is important to note that the operationalization of SES tends to vary across studies and sometimes is comprised of only one SES variable [60]. Such limitations can lead to difficulties in compiling and comparing results across studies. Importantly, although race and SES are

commonly correlated in health research, each factor has been found to relate somewhat differently to sleep processes and behaviors [61,62]. And therefore should be studied separately in relation to sleep loss and morbidity [63].

Adolescents of color

The demographic makeup of the nation has changed over the last 50 years. Today, the demographics of the nation's youth have transitioned to a non-White majority and will only continue to grow more diverse [64]. These changes provide a new sense of urgency and responsibility to conduct research using diverse subject populations¹. As an example, studies show that there are racial/ethnic differences in sleep processes and behaviors [e.g., 76-80]. Studies with African-American subject populations reveal that African-American youth have shorter total sleep times, higher levels of disrupted sleep, and more reported sleep problems than non-Hispanic White youth, even after controlling for SES [62,65,66]. There are also racial/ethnic differences in the daily levels of electronic media consumption [16]. And as described in a previous section, differences in the rates of obesity, all of which serve as risk factors for sleep problems. From this, we emphasize that all future research will need to have an intentional focus on including populations, particularly youth populations, from diverse racial, ethnic, and cultural backgrounds.

Socioeconomic environment

Residence within socioeconomically disadvantaged home environments early in life is associated with emotional, social, and biological hardships, and predicts future negative health outcomes [67,68]. Relatedly, there also is a significant association between low SES and poor sleep, in that adolescents living in disadvantaged neighborhoods have a greater incidence of adverse sleep behaviors (e.g., insufficient sleep) [69,70]. Specifically, individuals from low SES backgrounds report higher rates of sleep disturbance, problematic bed time behaviors, shorter total sleep time, and poorer quality sleep [61,71,72].

Outcomes of poor sleep have been associated with a factor found in low SES environments. For example, low-income or urban housing accommodations are often overcrowded or have thin walls separating the living quarters. This may increase residents' exposure to environmental

noise, and thereby contribute to shortened or fragmented sleep. In addition, pervasive environmental light exposure (e.g., street lamps) has demonstrated effects on circadian processes and contribute to adverse sleep behavior [71,73]. And there is also some evidence to suggest that low-income neighborhoods stand to be less safe, both inside and outside of the home [74,75]. Thus, living, working, and playing in certain environments may create elevated levels of threat and distress in certain youth populations. Such chronic adversity can activate biological stress responses (e.g., release of epinephrine, cortisol), promote nighttime arousal, and contribute to difficulties in falling or staying asleep [75].

Sleep behaviors of adolescents in rural communities should also be considered. In certain districts, many children commute, often by bus, to and from their school locations for up to 4 hours a day [76,77]. The burdens of early school start times and long commutes can result in significant reduction in overall sleep time [10,78]. External factors such as these may have cumulative effects on sleep patterns and short- and long-term health outcomes; however, the precise timing and duration of exposure necessary to produce these effects remain unknown.

Conclusions

There are complex social, cultural, and environmental factors that shape the quality and quantity of adolescent sleep and have the potential to create a lasting impact on the health outcomes of our nation. For the sake of brevity, this commentary did not present an exhaustive list of risk factors for sub-optimal sleep, nor was there an extensive review of the overlapping relations among the factors—although there would be value in doing so in future publications.

References

1. Centers for Disease Control and Prevention (2018) Sleep and Sleep Disorders. CDC Centers for Disease Control and Prevention.
2. Roth T, Ancoli-Israel S (1999) Daytime consequences and correlates of insomnia in the United States: results of the 1991 National Sleep Foundation Survey. II. *Sleep* 22(Suppl 2): S354-S358.
3. Benca RM, Obermeyer WH, Thisted RA, Gillin JC (1992) Sleep and psychiatric disorders. A meta-analysis. *Arch Gen Psychiatry* 49(8): 651-668.
4. Foster RG, Wulff K (2005) The rhythm of rest and excess. *Nat Rev Neurosci* 6(5): 407-414.

¹It is important to note that race/ethnicity may serve as a marker, but not as an explanation of group differences. That is, it is not yet clear as to whether the risk for suboptimal sleep in adolescents is due to a specific ethnic identity, a unique set of cultural influences, or some other variable.

5. Colten HR, Altevogt BM (2006) Sleep disorders and sleep deprivation: an unmet public health problem. National Academies Press. Washington DC, USA.
6. Hauri P, Olmstead E (1980) Childhood-onset insomnia. *Sleep* 3(1): 59-65.
7. Carskadon MA, Dement WC (2017) Normal human sleep: an overview. In: Kryger M, et al. (Eds.), *Principles and practice of sleep medicine*, p. 13-23.
8. Carskadon MA (1990) Patterns of sleep and sleepiness in adolescents. *Pediatrician* 17(1): 5-12.
9. Shochat T, Cohen-Zion M, Tzischinsky O (2014) Functional consequences of inadequate sleep in adolescents: A systematic review. *Sleep Med Rev* 18(1): 75-87.
10. Parthasarathy S, Carskadon MA, Jean-Louis G, Owens J, Bramoweth A, et al. (2016). Implementation of Sleep and Circadian Science: Recommendations from the Sleep Research Society and National Institutes of Health Workshop. *Sleep* 39(12): 2061-2075.
11. Moore M, Meltzer LJ (2008) the sleepy adolescent: causes and consequences of sleepiness in teens. *Paediatr Respir Rev* 9(2): 114-120.
12. Wittmann M, Dinich J, Merrow M, Roenneberg T (2006) Social jetlag: misalignment of biological and social time. *Chronobiol Int* 23(1-2): 497-509.
13. Dahl RE (2006) Sleeplessness and aggression in youth. *J Adolesc Health* 38(6): 641-642.
14. Wheaton AG, Jones SE, Cooper AC, Croft JB (2018) Short Sleep Duration among Middle School and High School Students-United States, 2015. *Morb Mortal Wkly Rep* 67(3): 85-90.
15. Calamaro CJ, Mason TB, Ratcliffe SJ (2009) Adolescents living the 24/7 lifestyle: effects of caffeine and technology on sleep duration and daytime functioning. *Pediatrics* 123(6): 1005-1010.
16. Rideout JV, Foehr UG, Roberts DF (2010) Generation M2: Media in the Lives of 8-18 year olds. A Kaiser Family Foundation Study.
17. Cain N, Gradisar M (2010) Electronic media use and sleep in school-aged children and adolescents: A review. *Sleep Med* 11(8): 735-742.
18. Hale L, Guan S (2015) Screen time and sleep among school-aged children and adolescents: a systematic literature review. *Sleep Med Rev* 21: 50-58.
19. Cajochen C, Frey S, Anders D, Spati J, Bues M, et al. (2011). Evening exposure to a light-emitting diodes (LED)-backlit computer screen affects circadian physiology and cognitive performance. *J Appl Physiol* 110(5): 1432-1438.
20. Wood B, Mark SR, Barbara P, Mariana GF (2013) Light level and duration of exposure determine the impact of self-luminous tablets on melatonin suppression. *Applied Ergonomics* 44(2): 237-240.
21. Lockley SW, Brainard GC, Czeisler CA (2003) High sensitivity of the human circadian melatonin rhythm to resetting by short wavelength light. *J Clin Endocrinol Metab* 88(9): 4502-4505.
22. Thapan K, Arendt J, Skene DJ (2001) An action spectrum for melatonin suppression: evidence for a novel non-rod, non-cone photoreceptor system in humans. *J physiol* 535(1): 261-267.
23. West KE, Jablonski MR, Warfield B, Cecil KS, James M, et al. (1985). Blue light from light-emitting diodes elicits a dose-dependent suppression of melatonin in humans. *J Appl Physiol* 110(3): 619-626.
24. Scheer FA, van Doornen LJ, Buijs RM (1999) light and diurnal cycle affect human heart rate: possible role for the circadian pacemaker. *J Biol Rhythms* 14(3): 202-212.
25. Cajochen C, Munch M, Koblalka S, Krauchi K, Steiner R, et al. (2005) High sensitivity of human melatonin, alertness, thermoregulation, and heart rate to short wavelength light. *J Clin Endocrinol Metab* 90(3): 1311-1316.
26. Noland H, Price JH, Dake J, Telljohann SK (2009) Adolescents' sleep behaviors and perceptions of sleep. *J Sch Health* 79(5): 224-230.
27. Orbeta RL, Overpeck MD, Ramcharran D, Kogan MD, Ledsky R (2006) High caffeine intake in adolescents: associations with difficulty sleeping and feeling tired in the morning. *J Adolesc Health* 38(4): 451-453.
28. Kaminer Y (2010) Problematic use of energy drinks by adolescents. *Child Adolesc Psychiatr Clin N Am* 19(3): 643-650.

29. Fisone G, Borgkvist A, Usiello A (2004) Caffeine as a psychomotor stimulant: mechanism of action. *Cell Mol Life Sci* 61(7-8): 857-872.
30. Karacan I, Thornby JI, Anch M, Booth GH, Williams RL, et al. (1976). Dose-related sleep disturbances induced by coffee and caffeine. *Clin Pharmacol Ther* 20(6): 682-689.
31. National Center for Chronic Disease Prevention and Health Promotion (2012) Preventing tobacco use among youth and young adults: A report of the Surgeon General. Centers for Disease Control and Prevention.
32. David LE, Leslie YK, Stratton K (2018) Public health consequences of e-cigarettes. The National Academies Press, Washington DC, USA.
33. Wetter DW, Young TB (1994) The relation between cigarette smoking and sleep disturbance. *Prev Med* 23(3): 328-334.
34. Jaehne A, Unbehau T, Feige B, Lutz UC, Batra A, et al. (2012). How smoking affects sleep: a polysomnographical analysis. *Sleep Med* 13(10): 1286-1292.
35. Benowitz NL (2008) Neurobiology of nicotine addiction: implications for smoking cessation treatment. *The Am J Med* 121(4): S3-S10.
36. Zhang L, Samet J, Caffo B, Punjabi NM (2006) Cigarette smoking and nocturnal sleep architecture. *Am J Epidemiol* 164(6): 529-537.
37. Gillin JC, Lardon M, Ruiz C, Golshan S, Salin-Pascual R (1994) Dose-dependent effects of transdermal nicotine on early morning awakening and rapid eye movement sleep time in nonsmoking normal volunteers. *J Clin Psychopharmacol* 14(4): 264-267.
38. Davila DG, Hurt RD, Offord KP, Harris CD, Shepard JW Jr (1994) acute effects of transdermal nicotine on sleep architecture, snoring, and sleep---disordered breathing in nonsmokers. *Am J Respir Crit Care Med* 150(2): 469-474.
39. Staner L, Luthringer R, Dupont C, Aubin HJ, Lagrue G (2006) Sleep effects of a 24-h versus a 16-h nicotine patch: a polysomnographic study during smoking cessation. *Sleep Med* 7(2): 147-154.
40. Hughes JR (2007) Effects of abstinence from tobacco: valid symptoms and time course. *Nicotine Tob Res* 9(3): 315-327.
41. Irwin CE Jr, Millstein SG (1986) Biopsychosocial correlates of risk-taking behaviors during adolescence. Can the physician intervene? *J Adolesc Health Care* 7(6 Suppl): 82S-96S.
42. Centers for Disease Control and Prevention (2008) Alcohol-related disease impact (ARDI). CDC, Atlanta, USA.
43. Johnson EO, Roehrs T, Roth T, Breslau N (1998) Epidemiology of alcohol and medication as aids to sleep in early adulthood. *Sleep* 21(2): 178-186.
44. Miller NS (2012) The pharmacology of alcohol and drugs of abuse and addiction. Springer
45. Ebrahim IO, Shapiro CM, Williams AJ, Fenwick PB (2013) Alcohol and sleep I: effects on normal sleep. *Alcohol Clin and Exp Res* 37(4): 539-549.
46. Roehrs T, Roth T (2001) Sleep, sleepiness, sleep disorders and alcohol use and abuse. *Sleep med Rev* 5(4): 287-297.
47. Robert Wood Johnson Foundation (2018) The state of Childhood obesity. The State of Obesity.
48. Chen X, Beydoun MA, Wang Y (2008) Is Sleep Duration Associated With Childhood Obesity? A Systematic Review and Meta-analysis. *Obesity (Silver Spring)* 16(2): 265-274.
49. Lytle LA, Pasch KE, Farbakhsh K (2011) The relationship between sleep and weight in a sample of adolescents. *Obesity (Silver Spring)* 19(2): 324-331.
50. Knutson KL, Spiegel K, Penev P, Van Cauter E (2007) The metabolic consequences of sleep deprivation. *Sleep Med Rev* 11(3): 163-178.
51. Spiegel K, Leproult R, Van Cauter E (1999) Impact of sleep debt on metabolic and endocrine function. *Lancet* 354(9188): 1435-1439.
52. Leproult R, Van Cauter E (2010) Role of sleep and sleep loss in hormonal release and metabolism. *Endocr Dev* 17: 11-21.
53. Spiegel K, Tasali E, Penev P, Van Cauter E (2004) Brief communication: sleep curtailment in healthy young men is associated with decreased leptin levels, elevated ghrelin levels, and increased hunger and appetite. *Ann Inter med* 141(11): 846-850.
54. Spiegel K, Knutson K, Leproult R, Tasali E, Van Cauter E (2005) Sleep loss: a novel risk factor for insulin

- resistance and Type 2 diabetes. *J appl physiol* 99(5): 2008-2019.
55. Taheri S (2006) the link between short sleep duration and obesity: we should recommend more sleep to prevent obesity. *Arch Dis Child* 91(11): 881-884.
56. Gangwisch JE, Malaspina D, Boden-Albala B, Heymsfield SB (2005) inadequate sleep as a risk factor for obesity: analyses of the NHANES. *Sleep* 28(10): 1289-1296.
57. Lopata M, Onal E (1982) Mass loading, sleep apnea, and the pathogenesis of obesity hypoventilation. *Am Rev Respir Dis* 126(4): 640-645.
58. Marshall NS, Glozier N, Grunstein RR (2008) Is sleep duration related to obesity? A critical review of the epidemiological evidence. *Sleep med Rev* 12(4): 289-298.
59. Peppard PE, Young T, Palta M, Dempsey J (2000) Skatrud J Longitudinal study of moderate weight change and sleep---disordered breathing. *JAMA* 284(23): 3015-3021.
60. Williams DR, Takeuchi DT, Adair RK (1992) Socioeconomic status and psychiatric disorder among blacks and whites. *Social Forces* 71(1):179-194.
61. El-Sheikh M, Kelly RJ, Buckhalt JA, Benjamin Hinnant J (2010) Children's sleep and adjustment over time: the role of socioeconomic context. *Child Dev* 81(3): 870-883.
62. Rao U, Poland RE, Lutchmansingh P, Ott GE, McCracken JT, et al. (1999) Relationship between ethnicity and sleep patterns in normal controls: implications for psychopathology and treatment. *J Psychiatr Res* 33(5): 419-426.
63. Williams NJ, Grandne MA, Snipes A, Rogers A, Williams O, et al. (2015) Racial/ethnic disparities in sleep health and health care: importance of the sociocultural context. *Sleep Health* 1(1): 28-35.
64. Jonathan V, David M Armstrong, Lauren M (2018) Demographic turning points for the united states: Population projections for 2020 to 2060. US Census Bureau.
65. Profant J, Ancoli-Israel S, Dimsdale JE (2002) Are there ethnic differences in sleep architecture? *Am J Hum Biol* 14(3): 321-326.
66. Nunes J, Jean-Louis G, Zizi F, Casimir GJ, von Gizycki H, et al. (2008) Sleep duration among black and white Americans: results of the National Health Interview Survey. *J Natl Med Assoc* 100(3): 317-322.
67. Leventhal T, Brooks-Gunn J (2000) the neighborhoods they live in: the effects of neighborhood residence on child and adolescent outcomes. *Psychol Bull* 126(2): 309-337.
68. Shonkoff JP, Phillips DA (2000) from neurons to neighborhoods: The science of early childhood development. National Academies Press, Washington DC, US.
69. Marco CA, Wolfson AR, Sparling M, Azuaje A (2011) Family socioeconomic status and sleep patterns of young adolescents. *Behav Sleep Med* 10(1): 70-80.
70. Tomfohr LM, Ancoli-Israel S, Dimsdale JE (2010) Childhood socioeconomic status and race are associated with adult sleep. *Behav Sleep Med* 8(4): 219-230.
71. Hill, TD, Burdette AM, Hale L (2009) Neighborhood disorder, sleep quality, and psychological distress: testing a model of structural amplification. *Health Place* 15(4): 1006-13.
72. Sichel AE, Moore PJ, Adler NE, Williams DR, Jackson JS (1999) The differential effects of sleep quality and quantity on the relationship between SES and health. *Ann N Y Acad Sci* 896: 431-434.
73. Dahl RE, Lewin DS (2002) Pathways to adolescent health sleep regulation and behavior. *J Adolesc Health* 31(6 Suppl): 175-184.
74. Brooks Gunn J, Duncan GJ, Aber JL (1997) Neighborhood poverty Volume 1: Context and consequences for children. Russell Sage Foundation, pp. 356.
75. Steptoe A, Feldman PJ (2001) Neighborhood problems as sources of chronic stress: development of a measure of neighborhood problems, and associations with socioeconomic status and health. *Ann Behav Med* 23(3): 177-185.
76. Pereira ÉF, Moreno C, Louzada FM (2014) Increased commuting to school time reduces sleep duration in adolescents. *Chronobiol Int* 31(1): 87-94.
77. Zars B, Long Rides, Tough Hides (1998) Long Rides, Tough Hides: Enduring Long School Bus Rides. ERIC.

78. Fox M (1996) Rural School Transportation as a Daily Constraint in Students' Lives. *Rural Educator* 17(2): 22-27.
79. Surani S, Subramanian S (2011) *Sleep and Safety*. Bentham Science Publishers
80. Mullington JM, Haack M, Toth M, Serrador JM, Meier-Ewert HK (2009) Cardiovascular, inflammatory, and metabolic consequences of sleep deprivation. *Prog in cardiovasc Dis* 51(4): 294-302.
81. McEwen BS (2006) Sleep deprivation as a neurobiologic and physiologic stressor: Allostasis and allostatic load. *Metabolism* 55(10Suppl2): S20-3.
82. Tsuno N, Besset A, Ritchie K (2005) Sleep and depression. *J Clin Psychiatry* 66(10): 1254-1269.
83. Harvey AG, Murray G, Chandler RA, Soehner A (2011) Sleep disturbance as trans diagnostic: consideration of neurobiological mechanisms. *Clin Psychol Rev* 31(2): 225-235.
84. Laberge L1, Petit D, Simard C, Vitaro F, Tremblay RE, et al. (2001). Development of sleep patterns in early adolescence. *J Sleep Res* 10(1): 59-67.
85. Carskadon MA, Wolfson AR, Acebo C, Tzischinsky O, Seifer R (1998) Adolescent sleep patterns, circadian timing, and sleepiness at a transition to early school days. *Sleep* 21(8): 871-881.
86. Arnett JJ (1999) Adolescent storm and stress, reconsidered. *Am Psychol* 54(5): 317-326.
87. Carskadon MA, Vieira C, Acebo C (1993) Association between puberty and delayed phase preference. *Sleep* 16(3): 258-262.
88. Roane BM, Taylor DJ (2008) Adolescent insomnia as a risk factor for early adult depression and substance abuse. *Sleep* 31(10): 1351-1356.
89. National Sleep Foundation (NSF) (2006) *Sleep in America Poll: Teens and sleep*. National Sleep Foundation.
90. Badia P, Myers B, Boecker M, Culpepper J, Harsh JR (1991) Bright light effects on body temperature, alertness, EEG and behavior. *Physiolbehave* 50(3): 583-588.
91. Mezick EJ, Matthews KA, Hall M, Strollo PJ Jr, Buysse DJ, et al. (2008) Influence of race and socioeconomic status on sleep: Pittsburgh Sleep SCORE project. *Psychosom Med* 70(4): 410-416.
92. Cooper RS, Kaufman JS, Ward R (2003) Race and genomics. *N Engl J Med* 348(12):1166-1170.
93. Hale L, Do DP (2007) Racial differences in self-reports of sleep duration in a population-based study. *Sleep* 30(9): 1096-103.
94. Combs D, Goodwin JL, Quan SF, Morgan WJ, Parthasarathy S (2016) Longitudinal differences in sleep duration in Hispanic and Caucasian children. *Sleep med* 18: 61-66.
95. Troxel WM, Ewing B, D'Amico EJ (2015) Examining racial/ethnic disparities in the association between adolescent sleep and alcohol or marijuana use. *Sleep Health* 1(2):104-108.
96. Chen E (2004) Why Socioeconomic Status Affects the Health of Children A Psychosocial Perspective. *Current Directions in Psychological Science* 13(3): 112-115.
97. Durkin MS, Davidson LL, Kuhn L, O Connor P, Barlow B (1994) Low-income neighborhoods and the risk of severe pediatric injury: a small-area analysis in northern Manhattan. *Am J Public Health* 84(4): 587-592.