

Effect of Ascorbic Acid on the Performance and Carcass Characteristics of Broiler Birds in Semi-Arid Zone, Sokoto State, Nigeria

Abdullahi AU^{1*}, Usman IS¹, Jafaru Y¹, Bello A² and Aliyu S³

¹Department of Animal science, Faculty of Agriculture, Usmanu Danfodiyo University, Nigeria

²College of Agriculture and Animal Science, Nigeria

³Department of Veterinary Anatomy, Faculty of Veterinary Sciences, Usmanu Danfodiyo University, Nigeria

*Corresponding author: Abdullahi AU, Department of Animal science, Faculty of Agriculture, Usmanu Danfodiyo University, Sokoto, Nigeria, Tel: +234(0)8039687589; Email: abccrcfge28@gmail.com

Received Date: April 15, 2020; Published Date: May 15, 2020

Abstract

Marshal strain of broiler chicken were used to determine the effect of Ascorbic acid (AA) on the performance and carcass characteristics of broiler birds at starter and finisher phase in Sokoto, Nigeria at different levels of AA inclusion. 225 day-old chicks were housed in a standard open sided tropical house type under deep litter system. The birds were randomly assigned to three treatments (0g, 10g and 20gAA) replicated Five times in completely randomize design (CRD). Feed intake (FI), body weight (BW), body weight gain (BWG) and feed conversion ratio (FCR) were determined. At the end of the experiment, 15 birds were randomly selected from each treatment, fasted, killed and their carcass weight analyzed. The result showed that there is significance difference $P < (0.05)$ between BW and FCR at starter phase. And no significant difference $P > (0.05)$ between FI and BWG. At finisher phase, the result showed that there is significance difference $P < (0.05)$ between BW and BWG and significance difference $P > (0.05)$ does not exist between FI and FCR across all the treatments. Birds fed diet with 0g AA performed better compared to other treatments in terms of FCR at finisher phase. While in the finisher phase those fed with 10g AA performed better compared to other treatments on the remaining parameters and also better in FCR at starter phase. Carcass analysis revealed significance difference $P < (0.05)$ between the treatment means with response to drumstick, legs, neck and gizzard while the other cuts and organs were statistically similar. Birds fed with 10g were superior in most of the parameters mentioned. It's recommended that to include AA in broiler diets at 10g level of AA.

Keywords: Broiler chicken; Ascorbic acid; Carcass characteristics; Body weight; Performance

Abbreviations: AA: Ascorbic acid; CRD: completely randomize design; FI: Feed intake; BW: body weight; BWG: body weight gain; FCR: feed conversion ratio; NRC: National Research Council; LSD: Least Significant Difference; TFI: Total feed intake; AFIB: Average feed intake per bird; AFIBD: Average feed intake per bird per day; TBW: Total body weight; BWB: Body weight per bird; BWBD: Body weight per

bird per day; BWGB: Body weight gain per bird; MWFI: Mean weekly feed intake.

Introduction

Poultry is a general term for birds of several species such as chicken or domestic fowls, ducks, geese, guinea fowls,

pigeons, ostriches and other game birds. In Nigeria, different poultry species contribute significantly to the animal protein supply of the populace in terms of eggs laid and meat produced. The general objective of poultry nutrition is to maximize the production performance of birds. Profitability in the poultry industry depends largely on the biological efficiency of the birds in terms of high egg production, efficiency of feed consumption and life ability. The economic efficiency of chicken meat appears to be dependent on the growth rate of the birds [1-4]. Over the years, poultry industry, chicken in particular has been the major source of animal protein in Nigeria. A substantial proportion of Nigeria foreign exchange goes to the importation of livestock, day old chicks and other foreign inputs in order to satisfy the local demand. However, the massive importation of poultry meat cannot provide a lasting solution to the problem of increasing animal poultry demand in Nigeria. Therefore, a planned development of livestock industry and increased poultry production would appear to be a better way of closing the protein gap at both short and long demand. Increase in egg and meat production can be achieved through proper nutrition. Feed is an important aspect of poultry production. Poultry production especially broiler chicken remains one of the viable way of achieving sustainable and rapid production of high quality animal to meet the increasing demand of the Nigerian teeming population. However, high cost of energy and protein concentrates used in formulation of poultry feeds has remained a significant challenge to the poultry industry [5-10].

Ascorbic acid (AA) also known as Vitamin C produces anti-stress hormones which help to combat the effect of stress caused on animal body. AA is a water soluble anti-oxidant that is normally produced in adequate quantity by metabolism in birds when not stressed. Butcher and Miles explained that its use as an anti-stress agent is often considered during periods of heat-stress since AA plays a major role in the biosynthesis of corticosterone, a hormone that enhances energy supply during stress. Commercial broilers are economic agricultural field production units in which the objective is to maximize field performance. Poultry require supplementary dietary vitamins since common feed ingredients used in poultry production do not provide adequate quantities to meet minimum requirements. Broilers are in continuous stress due to faster growth rate, pathogens, and the ever-changing environmental conditions in the broiler houses. Poultry have the ability to synthesize ascorbic acid, or vitamin C (VC), in their body; hence, no recommended requirement is established by the NRC [11-24]. That is why small scale broiler farmers are having difficulty in given the required dosage for the effective management of broilers. However, the study will help poultry farmers of the study area in solving the problems of effective use of supplemented ascorbic acid.

Poultry production has long been recognized as one of the quickest ways for rapid increase in protein supply in the shortest run, the study is carried out in the semi-arid zone, the area is characterized with higher temperature which is one of the constraint in poultry production, Ascorbic acid (AA) also known as Vitamin C produces anti-stress hormones which help to combat the effect of stress caused on animal body, the study will help poultry farmers in the effective use of supplemented ascorbic acid to reduce the effect of heat stress. The main objective of the study is to assess the effect of ascorbic acid on the performance and carcass characteristics of broiler birds in semi-arid zone Sokoto, Nigeria the specific objectives are:

- I. To assess the effect of ascorbic acid on the performance of broiler birds
- II. To assess the effect of ascorbic acid on the carcass characteristics of broiler birds
- III. To evaluate the performance and carcass characteristic at different level of Ascorbic acid inclusion

Materials and Methods

Description of the Study Area

The experiment was conducted in the poultry production and research unit of the Department of Animal Science, Faculty of Agriculture, Usmanu Danfodiyo University, Sokoto, at Sokoto State veterinary Centre, Aliyu Jedo road in Sokoto metropolis. Sokoto State is located between latitudes 13^o 05 and 13^o08'N and longitudes 5^o15' and 5^o25'E, within the Sudan savannah zone, in the extreme north western part of Nigeria and at an altitude of 350m above the sea level. It is bordered in the north with Niger Republic, Zamfara state to the east and Kebbi state to the south and west (SSMIYSC, 2010). The annual rainfall ranges between 500mm to 1,300mm. There are two major seasons in the state namely: wet and dry seasons. The dry season starts from October and last up to April, in some part and may extend to May to June in other part. The wet season on the other hand begins in most part of the state in May and last up to September or October [25,26].

Experimental Birds and their Management

Two hundred and twenty-five (225) day old broiler chicks (Marshall) was used for the study. The chicks were housed in a standard open-sided tropical house type under deep litter system. The housing was disinfected and fumigated, as well as thorough cleaning. The drinkers and feeders will also be kept clean, water and feed was provided *ad libitum*. The floor was covered using litter such as wood shavings. Administration of vaccine was accordingly and disease outbreak was managed adequately

Experimental Feed Material and their Sourcing

Commercial feed was given for 4 days; feed was self-formulated without the test ingredient which was given for 4 days which served as adjustment; afterward the test ingredient (Ascorbic acid) was added to the self-formulated feed which was given for 7 weeks. The feed ingredients which include maize, soya bean, groundnut cake, wheal offal, was sourced from Sokoto State central market. Other ingredients which include; bone meal, limestone, premix, salt, lysine and methionine was obtained from supply store.

Feed Formulation

The feed was self-formulated; the process involves in the formulation include.

Milling/Grinding

The size of the ingredient was reduced by grinding using a simple grinder which will help in the improvement of the

digestibility and result in more uniform feed.

Weighing of Ingredient

The grounded ingredient was weighed using weighing scale, some of the ingredients was weighed in grams (g) and others in kilograms (kg).

Mixing

Mixing was carried out on a clean floor, proper mixing of the grounded ingredients was ensured by the following step:

- Assembled ingredients was spread on the cleaned floor one after the other in a distinct layer
- Smaller ingredient was premixed with wheat-offal or ground grains before spreading
- Shoveling the feed from one end of the pile to the other to form a cone shaped pile,
- The mixture was shoveled in this manner for five (5) or more times to produce a homogenous mixture. The mixed feed was bagged in a dry bag for storage (Table 1).

Ingredients	Starter (Kg)	Finisher (Kg)
Feed composition		
Maize	50	49
Soya bean meal	18	22
Groundnut cake	20	12
Wheal offal	8	13
Limestone	1.5	0.5
Bone meal	1.5	2.5
Premix	0.25	0.25
Lysine	0.25	0.25
Methionine	0.25	0.25
Salt	0.25	0.25
Total	100	100
Composition	Starter diet (%)	Finisher diet (%)
ME (kcl/kg)	3054	2941
CP	23	21
Lysine	0.9	1.1
Methionine	0.6	0.6
Ca	1	0.9
P	0.5	0.6
Fibre	6	5.4

Table 1: Gross Feed and Chemical Composition of the Starter and Finisher Diet.

Experimental Design

Total number of birds was 225, Completely Randomize Design (CRD) was used, Three treatments (75 birds each)

and five replicates (15 birds each). Treatment one (T1) served as control with 0% ascorbic acid, Treatment two (T2) was containing 10g of Vitamin C in 10kg feed, Treatment

three (T3) was containing 20g of ascorbic acid in 10kg feed.

Data Collection

Performance

To assess the performance of the bird the following parameters was measure.

Feed Intake

Feed intake was recorded in daily basis by subtracting the weight of the remnant from weight of the quantity given the previous day.

Body Weight Gain

The weight gain was recorded on weekly basis by using a weighing scale.

Feed Conversion Ratio

The feed conversion ratio was determined by dividing average feed intake per bird by body weight per bird.

Mortality

Mortality was recorded on daily basis.

Carcass Characteristics

Primal cuts: Birds from different treatment was randomly selected for Primal cut, which was used to assess the carcass characteristics, the primal cut includes: the drumstick, thighs, breast, wings, bark, and neck.

Organs: Organs to be assess include gizzard, liver, heart, lungs, crop, spleen, head, intestine, bile, abdominal fat.

Weight and percentages: The percentage of live weight, killed weight, plucked weight (carcass), and dressed weight was recorded.

Data Analysis

The data collected was analyzed using descriptive statistics i.e. means percentages, ranges. Some data were analyzed using analysis of variance (ANOVA). Significant means was separated using Least Significant Difference (LSD).

Results and Discussion

Performance Characteristics at Starter Phase

The result on the performance characteristics of experimental broiler birds are shown in Table 2.

Parameters	T1(0g) Control	T1(10g)AA	T3(20g)AA	SEM*
Total feed intake (TFI)	14442.18	14128.07	14111.34	176.8181
Average feed intake per bird (AFIB)	1368.35	1284.37	1333.99	42.49953
Average feed intake per bird per day (AFIBD)	195.48	183.48	190.57	6.07136
Total body weight (TBW)	18155.72b	19092.30a	19677.67a	390.2168
Body weight per bird (BWB)	1717.88	1735.66	1862.3	63.58478
Body weight per bird per day (BWBD)	245.41	247.95	266.04	9.08354
Body weight gain (BWG)	7620.75	8141.87	8257.64	303.1631
Body weight gain per bird (BWGB)	719.3	740.17	781.8	30.6655
Body weight gain per bird per day (BWGBD)	102.76	105.74	111.69	4.38079
Feed conversion ratio	1.90b	1.74a	1.71a	0.06042

Table 2: The results on performance characteristic of broiler birds at starter phase.

Total Feed Intake (TFI)

Results of TFI performance are presented in Table 2 which shows that T1(control) has the highest mean feed intake of 14442.18g, followed by treatment T2(10g) with mean feed intake of 14128.07g while the last treatment T3 (20g AA) which has the lowest mean feed intake of 14111.34g at starter phase, with SEM of 176.1806 . The result indicates that there is no significant difference between the treatments. However the values are statistically the same but slightly differed numerically. Mean weekly feed intake (MWF) of the experimental birds as shown in Appendix

1 and Figure 1 increases from week 2 to week 4 in all the treatment, and ranges from the lowest value (3073.03g) recorded in treatment 2 (10gAA) at week 2 to the highest (6648.43g) in treatment 1 (0gAA) at week 4. This shows that with inclusion of AA in the feed of the birds tend to eat less; precisely it shows that birds consumed diets without AA inclusion perform better in terms of feed intake when compared to those with AA inclusion at starter phase. This is in agreement with the findings of kafri and cherry and kutlu and forbes [27,28], who reported that dietary supplemental AA tended to reduce performance of broiler chickens.




Figure 1: Average Feed Intake at Starter Phase.

Average Feed Intake Per Bird (AFB)

Results on average feed Intake are presented in Table 2 which of the experimental broiler birds showed that T_2 was highest with (1368.35) for birds consuming 0gAA then followed by birds in T_3 with (1333.99) consuming 20gAA and lowest was (1284.37) for birds in T_1 on 10gAA. Results showed that significant difference ($p>0.05$) does not exist between T_1 , T_2 and T_3 . SEM recorded under AFB was 42.49953.

Average Feed Intake Per Bird Per Day (AFBD)

Results on Average feed Intake per bird per day of the experimental broiler birds are presented in Table 2 which shows that T_3 was lowest (183.48) for birds consuming 10gAA and (190.57) for birds in T_3 consuming 20gAA the highest being 195.48 for birds in T_1 on 0gAA. However, significant difference ($p>0.05$) does not exist between treatments. SEM recorded under AFBD was 6.07136.

Body Weight

Total body weight (TBW): Results of TBW are presented in Figure 2 Which shows that significant difference ($p<0.05$) exist between T_2 , T_3 and T_1 , while T_2 and T_3 were statistically similar but they numerically differed. SEM recorded under TBW was (390.21682). Results on total body weight of the experimental broiler birds indicated that TBW (g) was highest (19677.67.) in T_3 for birds consuming 20gAA then followed by birds in T_2 with (19092.30) consuming 10gAA and lowest was (18155.72) for birds in T_1 on 0gAA, which recommend that AA should be included in a feed diet of broiler birds at a level of 20g due to its influence on increase body weight at starter phase. Mckee et al. [29], also demonstrated that 150 mg AA/kg feed significantly alleviated heat-related reductions in weight gain of birds of the same age, sex, and breed. Mean weekly body weight of the experimental birds increase at an increasing rate as they

aged, as it was shown in Appendix 3 Figure 2 which ranges from the lowest (3347.10g) in treatment 1 (0gAA) at week 2 to the highest (9795.60g) in treatment 3 (20gAA) at week 4.




Figure 2: Average Weekly Body Weight at Starter Phase.

Body weight per bird (BWB): Results on body weight per bird of the experimental broiler birds are presented in Table 2 which shows that BWB (g) was highest (1862.30) in T_3 for birds consuming 20gAA, the medium being (1735.66) for birds in T_2 consuming 10g-AA and the lowest was 1717.88 for birds in T_1 on 0gAA. However, significant difference ($p>0.05$) does not exist between treatments. SEM recorded under BWB was 63.58478.

Body weight per bird per day (BWBD): The result of BWBD are presented in Table 2 where BWBD values of 245.41, 247.95 and 266.04 were recorded for birds T_1 , T_2 and T_3 . Results on body weight per bird per day indicated that BWBD (g) was lowest with (245.41) for birds in T_2 consuming 0gAA then followed by 247.95 for birds in T_2 consuming 10gAA and highest was 266.04 for birds in T_3 on 20gAA. However, significant difference does not ($p>0.05$) exist between T_1 , T_2 and T_3 . SEM recorded under BWBD was 9.08354.

Body weight gain (BWG): Results of BWG are presented in Table 2 Significant difference ($p>0.05$) does not exist between the treatments. SEM recorded under BWG was 303.163.5. Results on body weight gain of the experimental broiler birds indicated that BWG (g) was lowest (7620.75) in T_1 for birds consuming 0gAA then followed by (8141.87) in T_2 for birds consuming 10gAA and highest was 8257.64 for birds in T_3 on 20gAA. This indicate that 20gAA has more effect than the other levels of ascorbic acid inclusion (10gAA and 0gAA) which means that T_3 perform better than the other treatments in terms of body weight gain at starter phase. Mean weekly body weight gain of the experimental birds increase at an increasing rate as they age as it was in the feed intake as shown in Appendix 5 and Figure 3.




Figure 3: Average Weekly Body Weight Gain at Starter Phase.

Body weight gain per bird (BWGB): Results on body weight gain per bird of the experimental broiler birds are presented in Table 2 which shows that BWGB (g) was lowest (719.30) in T_1 for birds consuming 0gAA then followed by 740.17 for birds in T_2 consuming 10gAA highest was 781.80 for birds in T_3 on 20gAA. However, significant difference ($p > 0.005$) does not exist between Treatments. SEM recorded under BWGB was (30.66550).

Body weight gain per bird per day (BWGBD): Result of BWGBD are presented in Table 2 which shows that BWGBD values of 102.76, 105.74 and 111.69 were recorded for birds on T_1 , T_2 and T_3 . Results on body weight gain per bird per day (BWGBD) indicated that BWGBD (g) was highest (111.69) in T_3 for birds consuming 20gAA then followed by (105.74) in T_2 for birds consuming 10gAA and lowest was 102.76 for birds in T_1 on 0gAA. However, significant difference ($p > 0.05$) does not exist between treatment. SE M recorded under BWGBD was 4.38079.

Feed conversion ratio (FCR): In Table 2, the results of FCR recorded SEM of 0.06042. Results on Feed Conversion Ratio of the experimental broiler birds narrated that FCR was highest (1.90) in T_1 for birds consuming 0gAA then followed by (1.74) for birds in T_2 consuming 10gAA and lowest was 1.71 for birds in T_3 on 20gAA. Results showed that FCR means of T_2 , T_3 were statistically similar; but they differed numerically. However, significant difference ($p < 0.05$) exist between the treatments T_2 , T_3 and T_1 , even though the difference is not much [30-34]. The mean weekly feed conversion ratio of the experimental birds ranges from the lowest (1.60) the most desirable value in treatment 3 (20gAA) at week 3, to the highest and least desirable value (2.04) in treatment 1 (0gAA) in week 4 as shown in Appendix 7 and Figure 4.




Figure 4: Average Weekly Feed Conversion Ratio at Starter Phase.

Performance Characteristics at Finisher Phase

Results obtained from the experiment at starter phase on performance characteristics of broiler birds are presented in Table 3.

Parameters	T1 (0g) Control	T2 (10g) AA	T3 (20g) AA	SEM
Total feed intake (TFI)	31214.32	32107.49	26430.18	2682.63
Average feed intake per bird (AFIB)	2952.69b	2918.86b	3936.81a	309.3
Average feed intake per bird per day (AFIBD)	421.81b	416.98b	562.40a	44.19
Total body weight (TBW)	58814.22ab	62832.73a	47692.67b	5975.56
Body weight per bird (BWB)	5555.97b	5712.07b	6889.8a	376.4
Body weight per bird per day (BWBD)	793.71b	816.01b	984.26a	53.77
Body weight gain (BWG)	14182.96a	16465.15a	5527.52b	2804.77
Body weight gain per bird (BWGB)	1344.96ab	1496.83a	388.09b	449.36
Body weight gain per bird per day (BWGBD)	192.14ab	213.83a	55.44b	64.19
Feed conversion ratio (FCR)	2.21	1.95	3.73	1.96

Table 3: The results on performance characteristic of experimental broiler birds at finisher phase.

Total feed intake (TFI): Results of TFI performance are showed in Table 3 which indicate that T1(control), T2(10g) and T3(20g) had (31214.32g), (32107.49g) and (26430g).18g respectively with SME (2682.63). From the result obtained, T2(10g) had the highest mean value followed by T1(control) for birds consuming (0g AA) and then T3 the lowest for birds consuming (20g AA) . The result indicates that there is no significant difference between the treatments. However the means were statistically similar but differed numerically. Result on weekly mean feed intake of the experimental birds at finisher phase showed that there is increase in feed intake from week 5 to 6 but reduce from week 6th to 7th as shown in Appendix 2 and Figure 5.




Figure 5: Average Feed Intake at Finisher Phase.

However, the weekly mean feed intake at finisher phases ranges from highest (12209.33) in T₂ (10gAA) at week 6 to the lowest (7354.86) in T₃ 20gAA at week 7th. However, there is decline from week 6th to week 7th in feed intake and also across treatment. The fluctuation in feed intake across the treatments may be attribute to disease incidence in treatment3 (20gAA) in the 7th week of the experiment [35,36]. reported on his experiment; Effect of ascorbic acid supplementation on broiler chicken stock at different densities in a humid tropical environment, that The total feed intake of broilers on optimal stocking density was similar ($p>0.05$) in feed intake of birds on double optimal stocking density supplemented with 300mg AA/litre of water. The least feed intake was however recorded for broilers on double optimal stocking density without AA supplementation.

Average Feed intake per bird (AFB): Results on average feed Intake per bird of the experimental broiler birds are presented in Table 3 which revealed that T₁, T₂ were statistically similar which shows that significant different ($p>0.05$) exist between (T₁,T₂) and T₃. However T₃ has the highest value (3936.81g) for birds consuming 20gAA then followed by birds in T₁ with (2952.69g) consuming 0g AA

and lowest was 2918.86g for birds in T₂ on 10g AA.. SEM recorded under AFB was (309.30).

Average Feed intake per bird per day (AFBD): Results on average feed Intake per bird per day of the experimental broiler birds are presented in Table 3 which revealed that T₁, T₂ were statistically similar and significant different ($p>0.05$) exist between them and T₃. However T₃ has the highest value (562.40g) for birds consuming 20gAA then followed by birds in T₁ with (421.81g) consuming 0g AA and lowest was(416.18g) for birds in T₂ on 10g AA.. SEM recorded under AFBD was (44.19).

Body weight

Total body weight (TBW): The results of TBW are presented in Table 3 Significant difference ($p>0.05$) exist between all the treatments (T₁, T₂ and T₃). SEM recorded under TBW was (5975.56).However, the results on total body weight of the experimental broiler birds indicated that TBW (g) was highest (62832.73g) in T₂ for birds consuming 10gAA then followed by birds in T₁ with (58814.22g) consuming 0gAA and lowest was (47692.67g) for birds in T₃ on 20gAA. This shows that consumed 10gAA performed better compared to those with (20g and 0g) of ascorbic acid inclusion. In appendix 4 and Figure 6 the result on mean weekly body weight gain of the experimental bird shows that the body weight gain increase as the age of the birds increases. However body weight fluctuate only across the treatment but progressively increases through the weeks. Which indicate an increase from week 5th to week 7th but decline in all the treatment. The mean ranges from the lowest (1417.0g) in treatment 1 (0gAA) at week 5 to the highest (26144.82g) in treatment 2 (10gAA) at week 7 Figure 6.




Figure 6: Average Weekly Body Weight at Finisher Phase.

Body weight per bird (BWB): Results on body weight per bird of the experimental broiler birds are presented in Table 3 which shows that body weight per birds (BWB) was highest (6889.8g) in T₃ for birds consumed 20gAA, the medium

being (5712.07g) for birds in T_2 consumed 10gAA and the lowest was 5555.97g for birds in T_1 on 0gAA. However, there is significant difference ($p < 0.05$) between the treatments, in which T_2 and T_1 were statistically similar. SEM recorded under BWB was 376.40.

Body weight per bird per day (BWBD): In Table 3, the result of body weight per bird per day revealed that BWBD (g) values of (793.71), (816.01) and (984.26) were recorded for birds T_1 , T_2 and T_3 respectively. Results on body weight per bird per day indicated that BWBD (g) was lowest with (793.71) for birds in T_1 consuming 0gAA then followed by 816.01 for birds in T_2 consuming 10gAA which was the moderate and highest was 984.26 for birds in T_3 on 20gAA. However, significant difference ($p > 0.05$) exist between T_2 , T_1 and T_3 in which T_2 , T_1 were statistically similar but differed numerically. SEM recorded under BWBD was (53.77).

Body weight gain (BWG): Results of BWG are presented in Table 3 which revealed Significant difference ($p > 0.05$) between treatments (T_1 , T_2 and T_3) in which T_2 , T_1 were statistically similar. SEM recorded under BWG was 2804.77. Results on body weight gain of the experimental broiler birds indicated that BWG (g) was lowest (5527.52) in T_3 for birds consuming 20gAA then followed by (14182.96) in T_1 for birds consuming 0gAA which was moderate and highest was (16465.15) for birds in T_2 on 10gAA. Weekly body weight gain of the experimental birds at finisher phase as shown in Appendix 6 and Figure 7 ranges from the lowest (1927.41g) in treatment 1 at week 7 to the highest (7187.71g) at the same treatment at week 6 which indicate that there was decline across treatments and weeks. The fluctuation in body weight gain across the weeks and treatments may be attribute to disease incidence in T_3 and T_2 at 7th week of the experiment in figure 7.




Figure 7: Average Weekly Body Weight Gain at Finisher Phase.

Body weight gain per bird (BWGB): Results on body weight gain per bird of the experimental broiler birds are presented

in Table 3 which shows that BWGB (g) was lowest (388.09) in T_3 for birds consuming 20gAA then followed by (1344.96) for birds in T_1 consuming 0gAA highest was (1496.83) for birds in T_2 on 10gAA. However, significant difference ($p < 0.05$) exist between all the treatments. SEM recorded under BWGB was (449.36).

Body weight gain per bird per day (BWGBD): Results of body weight gain per bird per day are presented in Table 3 BWGBD values of (192.14), (213.83) and (55.44) were recorded for birds on T_1 , T_2 and T_3 respectively. Results on body weight gain per bird per day (BWGBD) indicated that BWGBD (g) was highest (213.83) in T_2 for birds consuming 10gAA then followed by (196.14) in T_1 for birds consuming 0gAA and lowest was (55.44) for birds in T_3 on 20gAA. However, significant difference ($p > 0.05$) exist between all the treatments. SEM recorded under BWGBD was (64.19).

Feed conversion ratio (FCR): The results of FCR are presented in Table 3 which recorded SEM of 1.96. The results on Feed Conversion Ratio of the experimental broiler birds narrated that FCR was highest (3.73) in T_3 for birds consuming 20gAA then followed by (2.21) for birds in T_1 consuming 0gAA and lowest was 1.95 for birds in T_2 on 10gAA. Results showed that FCR means of T_2 , T_3 and T_3 are statistically similar; but they differ numerically. However, significant difference ($p > 0.05$) does not exist between the treatments. The weekly feed conversion ratio of the experimental birds ranges from the lowest and most desirable value (1.18g) in treatment 3 (20gAA) at week 7th to the highest and least desirable (5.89g) in treatment 1 at week 7th. Which indicate that there was decline across treatments and weeks? The experimental birds in treatment 3 have the best feed conversion ratio due to the high level of Ascorbic Acid inclusion while for those in treatment 1 with 0 inclusion of Ascorbic Acid have poor feed conversion ratio in Figure 8.




Figure 8: Average Weekly Feed Conversion Ratio at Finisher Phase.

Carcass Characteristics of Boiler Birds

experimental birds are presented in Table 4.

The results of the carcass characteristics obtained from the

Parameters	T1(0g)	T2(10g) AA	T3(20g) AA	SEM
Drumstick (DRM)	188.80b	208.35a	212.65a	3.14
Thighs (THI)	251.8	251.5	249.6	6.33
Breast (BRE)	718	663.3	650.75	31.41
Wing (WNG)	166.5	165.1	162.3	4.53
Back (BCK)	167.7	175.35	172.05	6.71
Neck (NCK)	70.30b	76.30a	75.75a	2.29
Total primal cuts (TPRM)	1563.1	1539.9	1523.1	47.83
Gizzard (GZZ)	49.10a	41.20b	39.20b	3.07
Liver & Bile (L/B)	49.5	50.6	48.55	2.89
Heart (HRT)	8	9.4	7.8	0.75
Lung (LNG)	8.30b	9.90b	12.15a	0.78
Crop (CRP)	10.4	21.5	14.65	5.76
Spleen (SPLN)	1.7	2.05	1.6	0.38
LEGS	64.10b	69.50b	74.05a	2.92
HEAD	43.4	44.25	45.2	1.15
Intestine (INT)	124.2	128.1	122.5	7.73
Abdominal Fats (ABDF)	20.4	22.85	24.05	4.09
Total Organs (TORG)	379.1	399.35	389.75	15.4
Live weight (LW)	2157.3	2168.35	2133	52.07
Kill weight (KW)	2090.5	2092	2070.15	50.52
Pluck weight (PW)	2017.9	2012.45	1997.9	48.43
Dress weight (DW)	1661.7	1631.7	1610.85	47.86
	72.43	71.01	71.38	0.84
Primal cuts as percentage live weight (P/LW)				
Primal cuts as percentage dress weight (P/DW)	94.05	94.36	94.56	0.32
Organs cuts as percentage live weight (O/LW)	17.61	18.42	18.29	0.81
Organs cuts as percentage dress weight (O/DW)	22.87	24.53	24.24	1.27

Table 4: The result of carcass characteristics of broiler birds.

Drumstick: The result of drumstick is presented in Table 4 and Figure 9. SEM recorded under drumstick weight was 3.14. Results on drumstick weight indicated that drumstick Weight (g) was highest (212.65) in T₃ followed by T₂ (208.35)

and lowest was 188.80 for birds in T₁. However, significant difference ($p < 0.05$) exist between T₂, T₃ and T₁. In which T₂ and T₃ are statistically similar but differ numerically.




Figure 9: Average Primal Cuts at Finisher.

Thigh: The result of thigh is presented in Table 4 and Figure 9. SEM recorded under thigh Weight was 6.33. Results on thigh weight indicated that thigh Weight (g) was highest (251.80) in T_1 followed by T_2 with (251.50) and lowest was (249.60) for birds in T_3 . However, no significant difference ($p > 0.05$) exist between T_1 , T_2 and T_3 .

Wing: The result of wing is presented in Table 4 and figure 9. Wing Weight values of (166.50), 165.10 and (162.300 were recorded for birds on T_1 , T_2 and T_3 . However, no significant difference ($p > 0.05$) exist between T_1 , T_2 and T_3 . SEM recorded under wing Weight was (4.53). Results on wing weight of the experimental broiler birds indicated that wing Weight (g) was highest (166.50) in T_1 followed by T_3 and lowest was 162.30 for birds in T_3 .

Back: The result of back is presented in Table 4 and Figure 9. Back Weight values of (167.70g), (175.35g) and (172.05g) were recorded for birds on T_1 , T_2 and T_3 . However, no significant difference ($p > 0.05$) exist between T_1 , T_2 and T_3 . SEM recorded under Back Weight was (6.71). Results on back weight indicated that Back Weight (g) was highest (175.35g) in T_2 and lowest was (167.70g) for birds in T_1 .

Neck: The result of neck are presented in Table 4 and Figure 9. SEM recorded under neck weight was 2.29. Results on neck weight indicated that neck Weight (g) was highest (76.30) in T_2 followed by T_3 (75.75) and lowest was 70.30 for birds in T_1 . However, significant difference ($p < 0.05$) exist between T_2 , T_3 and T_1 . In which T_2 and T_3 are statistically similar but differ numerically.

Total primal cuts: The results of total primal cuts are presented in Table 4 and Figure 9. TPRM values of (1563.10), (1539.90) and (1523.10) were recorded for birds on T_1 , T_2 and T_3 . However, no significant difference ($p > 0.05$) exist between T_1 , T_2 and T_3 . SEM recorded under TPRM Weight was 47.83. Results on TPRM weight of the experimental

broiler birds indicated that TPRM Weight (g) was highest (1563.10) in T_1 followed by T_2 with (1539.90) and lowest was ((1523.10) for birds in T_3 .

Organs

Gizzard: The result of gizzard are presented in Table 4 and Figure 10. Significant difference ($p < 0.05$) exist between T_2 , T_3 and T_1 were T_2 and T_3 are statistically similar. SEM recorded under gizzard Weight was 3.07. However, results on Gizzard weight of the experimental broiler birds indicated that gizzard Weight (g) was highest (49.10) in for birds in T_1 followed by T_2 (41.20) and lowest were (39.20) for birds in T_3 .




Figure 10: Average Carcass Organs at Finisher Phase.

Liver and Bile: The result of liver and bile are presented in Table 4 and Figure 10. Results on liver weight of the experimental broiler birds indicated that liver Weight (g) was highest (50.60) in T_2 followed by birds in T_1 with (49.50) and lowest was 48.55 for birds in T_3 . However, no significant difference ($p > 0.05$) exist between the treatments. SEM recorded under liver Weight was 2.89.

Heart: The result of heart is presented in Table 4 and Figure 10. Significant difference ($p > 0.05$) does exist between T_1 , T_2 and T_3 . SEM recorded under heart Weight was 0.75. However, results on heart weight indicated that heart Weight (g) was highest (9.40) in T_2 then followed by birds in T_1 with (8.00) and lowest was 7.80 for birds in T_3 .

Lung: SEM recorded under lung Weight was 0.78. Results on lung weight indicated that lung Weight (g) was highest (12.15) for birds in T_3 then followed by T_2 with (9.90) and lowest was 8.30 for birds in T_1 . However, significant difference ($p < 0.05$) exist between T_1 , T_2 and T_3 . Where T_1 and T_2 are statistically similar.

Crop: Results on crop weight of the experimental broiler birds are presented in Table 4 and Figure 10. Crop Weight (g) was highest (21.50) for birds in T_2 then followed by (14.65) in T_3 and lowest was 10.40 for birds in T_1 . However, significant difference ($p>0.05$) does not exist between T_1 , T_2 and T_3 . SEM recorded under crop Weight was 5.76.

Spleen: The result of drumstick are presented in Table 4 and Figure 10. No significant difference ($p>0.05$) exist between the treatments. SEM recorded under Spleen Weight was 0.38. Results on spleen weight of the experimental broiler birds indicated that spleen Weight (g) was highest (2.05) for birds in T_2 then followed by (1.70) in T_1 and lowest was 1.60 for birds in T_3 .

Legs: Results on leg weight of the experimental broiler birds are presented in Table 4 and Figure 10. Leg Weight (g) was highest (74.05) for birds in T_3 then followed by birds in T_2 with (69.50) and lowest was 64.10 for birds in T_1 . However, significant difference ($p<0.05$) exist between T_1 , T_2 and T_3 . Where T_1 and T_2 are statistically similar. SEM recorded under leg Weight was 2.92.

Head: The result of head are presented in Table 4 and Figure 10. SEM recorded under head Weight was 1.15. Results on head weight indicated that head Weight (g) was highest (45.20) for birds in T_3 then followed by T_2 with (44.25) and lowest was 43.40 for birds in T_1 . However, significant difference ($p>0.05$) does not exist between treatments.

Intestine: Results on intestine weight are presented in Table 4 and Figure 10. Intestine Weight (g) was highest (128.10) for birds in T_2 then followed by bird in T_1 with (124.20) and lowest was (122.50) for birds in T_3 . However, no significant difference ($p>0.05$) exist between the treatments. SEM recorded under Intestine Weight was 7.73 [37].

Abdominal fat: Results on abdominal fat weight of the experimental broiler birds are presented in Table 4 and Figure 10. Abdominal fat Weight (g) was highest (24.05) in T_3 followed by birds in T_2 with (22.85) and lowest was (20.40) for birds in T_1 . However, no significant difference ($p>0.05$) exist between the treatments. SEM recorded under liver Weight was 4.09.

Total organs: Results on total organs of the experimental broiler birds are presented in Table 4 and Figure 10. Total organ values of (379.10), (399.35) and (389.75) were recorded for birds on T_1 , T_2 and T_3 . However, no significant difference ($p>0.05$) exist between T_1 , T_2 and T_3 . SEM recorded under Total organ Weight was (915.40). Results on Total organ weight of the experimental broiler birds indicated that Total organ Weight (g) was highest (399.35) in T_2 followed by T_3 with (389.75) and lowest was (379.10) for birds in T_1

[38-40].

Percentages

Live weight (%): The result of live weight are recorded in table 4.3 and figure 11. Live Weight values of (2157.30), (2168.35) and (2133.00) were recorded for birds on T_1 , T_2 and T_3 . However, no significant difference ($p>0.05$) exist between T_1 , T_2 and T_3 . SEM recorded under Live Weight was 52.07 Results on live weight indicated that live Weight (g) was highest (2168.35) in T_2 then in T_1 with (2157.30) and lowest was (2133.00) for birds in T_3 .

Killed Weight (%): The result of kill weight are recorded in table 4.3 and figure 11. Significant difference ($p>0.05$) does exist between T_1 , T_2 and T_3 . SEM recorded under kill Weight was 50.52 However, results on kill weight indicated that kill Weight (g) was highest (2092.00) in T_2 then followed by birds in T_1 with (2090.50) and lowest was (2070.15) for birds in T_3 .

Plucked Weight (%): The result of pluck weight are recorded in table 4.3 and Figure 11. SEM recorded under plucked Weight was 48.43. Results on plucked weight indicated that plucked Weight (g) was highest (2017.90) for birds in T_1 then followed by T_2 with (2012.45) and lowest was 1997.90 for birds in T_3 . However, significant difference ($p>0.05$) does not exist between treatments.




Figure 11: Average Weights of Experimental Birds.

Dressed weight: The result of dress weight is recorded in Table 4 and Figure 11. No significant difference ($p>0.05$) exist between the treatments. SEM recorded under Dressed Weight was 47.86. Results on Dressed weight of the experimental broiler birds indicated that Dressed Weight (g) was highest (1661.70) for birds in T_1 followed by (1631.70)

for birds in T_2 and lowest was (1610.85) for birds in T_3 .

Primal cord as percentage of live weight P/LW (%): The results obtained on percentage as live weight are recorded in Table 4, which shows that significant difference ($p>0.05$) does exist between T_1, T_2 and T_3 . SEM recorded under heart Weight was 0.84. However, results on primal cord (%) live weight indicated that P/LW Weight (g) was highest (72.43) in T_1 then followed by birds in T_3 with (71.38) and lowest was 71.01 for birds in T_2 .

Primal cord as percentage of Dressed Weight P/DW (%): SEM recorded under P/DW was 0.32. Results on head weight indicated that P/DW Weight (g) was highest (94.56) for birds in T_3 then followed by T_2 with (94.36) and lowest was 94.05 for birds in T_1 . However, significant difference ($p>0.05$) does not exist between treatments.

Organs as percentage of live weight O/LW (%): Significant difference ($p>0.05$) does exist between T_1, T_2 and T_3 . SEM recorded under O/LW Weight was 0.81. However, results on O/LW weight indicated that O/LW Weight (g) was highest (18.42) in T_2 then followed by birds in T_3 with (18.29) and lowest was (17.61) for birds in T_1 .

Organs as percentage Dressed Weight O/DW (%): The results of organs as percentage dress weight are recorded in Table 4. Results O/DW values of (22.87), (24.53) and (24.24) were recorded for birds on T_1, T_2 and T_3 . However, no significant difference ($p>0.05$) exist between T_1, T_2 and T_3 . SEM recorded under O/DW Weight was 1.27. Results on O/DW weight indicated that O/DW Weight (g) was highest (24.53) in T_2 then in T_3 with (24.24) and lowest was (22.87) for birds in T_1 .

Conclusion and Recommendation

Conclusions

Base on the result obtained, AA has the positive effect on the performance and carcass characteristics on broiler birds at starter and finisher phase. The research therefore could be concluded that AA be included in the diet of broiler birds at starter and finisher phase at level of 10g in the feed formulation.

Recommendations

Further research should be conducted on higher levels of AA inclusion to establish the optimum inclusion rates. Also, further research need to be conducted to compare efficiency on other levels of AA inclusion apart from feed. (e.g. in drinking water). Study should be undertaken to determine at with level of inclusion of AA it serve best as anti-stress for the broiler birds in the study area. Research should also

be conducted on the broiler birds fed with 0g of AA, that perform even better than of higher levels of AA inclusion.

References

1. Abidin Z, Khatoon A (2013) Heat stress in poultry and the beneficial effects of ascorbic acid (vitamin C) supplementation during periods of heat stress. *World's Poultry Sci J* 69(1): 135-152.
2. Agri-Food Canada (2006) Poultry Marketplace profile of the Canadian Chicken Industry (2006) Chapter 3: Broiler and Hatching Egg Segment of the Industry.
3. Alabi RA, Alabi O F (2007) Effect of Economic Liberation on chicken Meat production in Nigeria. *Eur J Social Sci* 7(4): 58-61.
4. al-Taweil RN, Kassab A (1990) Effect of dietary vitamin C on ascites in broiler chicks. *Int J Vitam Nutr Res* 60(4): 366-371.
5. Barroeta AC, Baucells MD, Blanco Perez A, Calsamiglia S, Casals R, et al. (2012) Optimum vitamin nutrition in poultry breeders, in Optimum vitamin nutrition; in the production of quality animal foods. 5M Publishing, Benchmark House, 8 Smithy Wood Drives, Sheffield, S35 1QN, United Kingdom. pp: 1-385.
6. Chiang-Ting Chien, Wei-Tien Chang, Heui-Wen Chen, Tzung-Dau Wang, Shaw-Yih Liou, et al. (2004) Ascorbate Supplement Reduces Oxidative Stress in Dyslipidemic Patients Undergoing Apheresis. *Arterioscler Thromb Vas Bio* 24(6): 1111-1117.
7. Cunningham D (2004) Contract Broiler Production: Questions and Answer. The Poultry Site.
8. Curca D, Andronie V, Andronie IC, Pop A (2004) The influence of feed supplementation with ascorbic acid and sodium ascorbate on broilers under thermal stress. *Book of Abstract XXII World Poultry Congress of WPSA, Istanbul, Turkey* pp: 290.
9. Damerow G (1995) Storey's Guide to Raising Chickens: Care / Feeding / Facilities. Storey Publisher, Nigeria.
10. Adene Df, Ae Oguntade, (2006) The Structure and Importance of the Commercial and Village Based Poultry Industry In Nigeria. pp: 28-32.
11. Darre MJ (2006) The Importance of Poultry in Our Lives.
12. Ducan IJH, Hocking PM, Seawright E (1990) Sexual behavior and fertility in broiler breeder domestic fowl. *Applied Animal Behaviour Sci* 26(3): 201-213.

13. Elfic D, International Product Manager, Aviagen International (2012) A Brief History of Broiler Selection: How Chicken Became a Global Food Phenomenon in 50 Years.
14. Eriksson J, Larson G, Gunnarsson U, Bed'hom B, Tixier-Boichard M, et al. (2008) Identification of the yellow skin Gene Reveals a Hybrid Origin of the Domesticated Chicken. *Genetics. Plos Genetics* 4(2): e1000010.
15. FAO (1995) Food and Agricultural organization of the united Nation Quaternary Report, production Book 35. Rome Italy pp: 75-78.
16. FAO (2009) Food and Agricultural Organisation of the United Nation. Avian Influenza Control Programme in Indonesia.
17. FAO (2012) Food Outlook, Rome: Trade and Market Division of FAO.
18. Farooq M, Mian MA, Asghar A (2001) Factors affecting cost of production and net profit per broiler in the Subtropics. *Livestock Research for Rural Development* 13(1).
19. Garrigus WP (2007) Poultry Farming. *Encyclopaedia Britannica*.
20. Kutlu HR, JM Forbes (1993) Self-selection of ascorbic acid in coloured foods by heat-stressed broiler chicks. *Physiol Behav* 53(1): 103-110.
21. Kutlu HR (2001) Influences of wet feeding and supplementation with ascorbic acid on performance and carcass composition of broiler chicks exposed to a high ambient temperature. *Arch Tierernahr* 54(2): 127-139.
22. Leach RM Jr, Burdette JH (1985) The influence of ascorbic acid on the occurrence of tibial dyschondroplasia in young broiler chickens. *Poult Sci* 64(6): 1188-1191.
23. Leeson S, JD Summers (2001) Scott's Nutrition of the Chicken. 4th (Edn).
24. NRE (1994) Nutrient Requirements of Poultry. 9th (Edn), National Academy of Science, National Research Council, Washington pp: 176.
25. Consensus Study Report (1987) Vitamin Tolerance of Animals. National Academy of Sciences National Research Council, Washington DC, pp: 108.
26. Olivares M, Pizzaro F, Pineda O, Name JJ, Hertrampf E, et al. (1997) Milk inhibits and ascorbic acid favors ferrous bis-glycine chelate bioavailability in humans. *J Nutr* 127(7): 1407-1411.
27. Oluyemi JA, FA Roberts (2000) Poultry production in Warm Wet Climate. Macmillan Publisher.
28. Pardue SL, Thaxton JP, Brake J (1985) Influence of supplemental ascorbic acid on broiler performance following exposure to high environmental temperature. *Poult Sci* 64(7): 1334-1338.
29. Portsmouth J (1978) Practical Poultry keeping. 7th (Edn), Spur Publishing Company Ltd, England pp: 68-99.
30. Rennie JS, Fleming RH, McCormack HA, McCorquodale CC, Whitehead CC (1997) Studies on effects of nutritional factors on bone structure and osteoporosis in laying hens. *Br Poult Sci* 38(4): 417-4124.
31. Sainsbury MJ (2000) Poultry Health and Management. 4th (Edn), Blackwell Publication, London pp: 361.
32. Satterfield GH, MA Mosley, HC Gauger, AD Holmes, F Tripp (1940) Correlation of avian diseases and ascorbic acid content of chicken blood. *Poult Sci* 19: 337-344.
33. Settar P, Yalcin S, Turkmat L, Ozkan S, Cahanar A (1999) Season by genotype interaction related to broiler growth rate and heat tolerance. *Poult Sci* 78(10): 1353-1358.
34. Sies H, Stahl W (1995) Vitamins E and C, beta-carotene, and other carotenoids as antioxidants. *Am J Clin Nutr* 62(6): 1315S-1321S.
35. Takahashi K, Y Akiba, M Horiguchi (1991) Effects of supplemental ascorbic acid on performance, organ weight and plasma cholesterol concentration in broilers treated with propylthiouracil. *Brit Poul Sci* 32(3): 545-554.
36. Tee ES, Mohd Ismail N, Mohd Nasir A, Kahtijah I (1998) Nutrient composition of Malaysian foods. *FAO United Kingdom*.
37. USDA (2012) US Broiler Industry: Background Statistics and Information.
38. USDA (2020) Livestock and Poultry: World Markets and Trade. USDA.
39. Weber P, Bendich A, Schalch (1996) Ascorbic acid and human health—a review of recent data relevant to human requirements. *Int J Vit Nutr Res* 66(1): 19-30.
40. Zee JA, Carmichael L, Codere D, Poirier D, Fournier M (1991) Effect of storage conditions on the stability of vitamin C in various fruits and vegetables produced and consumed in Quebec. *J Food Compos Anal* 4(1): 77-86.